

METHOD OF UNDETERMINED COEFFICIENTS

Method for finding a particular solution $Y(t)$ of a constant coefficient linear inhomogeneous equation, i.e., one of the form

$$ay''(t) + by'(t) + cy(t) = g(t),$$

for certain special forms of the function g . To use the method you must know the roots of the characteristic equation

$$ar^2 + br + c = 0. \tag{*}$$

We use the following notation: $P_n(t)$ and $P_n^*(t)$ denote polynomials in t of degree n ; these are part of the *given* function g :

$$P_n(t) = a_0t^n + \cdots + a_{n-1}t + a_n, \quad P_n^*(t) = a_0^*t^n + \cdots + a_{n-1}^*t + a_n^*.$$

$Q_n(t)$ and $Q_n^*(t)$ also denote polynomials in t of degree n , but with *undetermined* coefficients; these are part of the solution $Y(t)$:

$$Q_n(t) = A_0t^n + \cdots + A_{n-1}t + A_n, \quad Q_n^*(t) = A_0^*t^n + \cdots + A_{n-1}^*t + A_n^*.$$

Given function $g(t)$	Form of $Y(t)$	Condition
$e^{\alpha t}$	$Ae^{\alpha t}$ $Ate^{\alpha t}$ $At^2e^{\alpha t}$	if α is not a root of (*) if α is a simple root of (*) if α is a double root of (*)
$P_n(t)$	$Q_n(t)$ $tQ_n(t)$ $t^2Q_n(t)$	if 0 is not a root of (*) if 0 is a simple root of (*) if 0 is a double root of (*)
$P_n(t)e^{\alpha t}$	$Q_n(t)t^s e^{\alpha t}$	if α is a root of (*) of order s
$ae^{\alpha t} \cos \beta t + be^{\alpha t} \sin \beta t$	$e^{\alpha t}(A \cos \beta t + B \sin \beta t)$	if $\alpha + i\beta$ is not a root of (*)
$e^{\alpha t}[P_n(t) \cos \beta t + P_n^*(t) \sin \beta t]$	$e^{\alpha t}[Q_n(t) \cos \beta t + Q_n^*(t) \sin \beta t]$ $te^{\alpha t}[Q_n(t) \cos \beta t + Q_n^*(t) \sin \beta t]$	if $\alpha + i\beta$ is not a root of (*) if $\alpha + i\beta$ is a simple root of (*)