

**Definition of limit.** It is useful at this point to review the definition of limit, extended to cover functions of several variables. For simplicity, we use only two variables.

We say that

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L \quad (*)$$

if we can force  $f(x, y)$  to be arbitrarily close to  $L$  (within an  $\epsilon$  given by *el Exigente*) by demanding that  $(x, y)$  be sufficiently close to  $(a, b)$  (at a distance no more than a  $\delta$  that we choose), independent of direction.

Note that it is the whole sentence (\*) that is defined. The use of the equal sign in that definition is justified by . . . .

**The main theorem.** Limits are unique. That is, given the function  $f$  and the point  $(a, b)$ , there is *at most one* choice of  $L$  satisfying (\*).

*Proof.* If not, let  $L_0$  and  $L_1$  be two different numbers that both satisfy (\*). The triangle inequality gives

$$|L_0 - L_1| \leq |L_0 - f(x, y)| + |f(x, y) - L_1| \quad (\dagger)$$

for any point  $(x, y)$  in the domain of  $f$ . If you persist in claiming that  $L_0$  and  $L_1$  both satisfy (\*), I calculate

$$d = |L_0 - L_1|.$$

To say that  $L_0 \neq L_1$  is to say that  $d > 0$  (strictly). Then I choose  $\epsilon = d/2$ . Then, if  $(x, y)$  is any point that you think is close enough to  $(a, b)$ , then both terms on the right side of ( $\dagger$ ) are less than  $d/2$ , which contradicts ( $\dagger$ ).

**Non-existence of limits.** The idea of this proof is used to identify cases in which limits do not exist. If there are two different values  $v_0$  and  $v_1$  such that  $f(x, y) = v_0$  for some points  $(x, y)$  arbitrarily close to  $(a, b)$  while  $f(x, y) = v_1$  for other points arbitrarily close to  $(a, b)$ , then it is not possible for  $f(x, y)$  at *all* points  $(x, y)$  close to  $(a, b)$  to be within  $|v_0 - v_1|/2$  of any single value.

**Continuity.** Almost every function that we know how to express has the property that, at points where the expression for the function can be evaluated, that value may be used for  $L$  in (\*). This property is called **continuity**. Continuity means that the value of a function at a point can be approximated by evaluating the function at a nearby point. This is what we do every time we trust our calculators to do much more than verify that  $1 + 1 = 2$ , so continuity is an abstraction of the idea of *computable*. The rules for building functions that we use are easily seen to preserve continuity, although complicated functions (for which  $\delta$  is small) may require extraordinary care to give reasonable accuracy.

### Exercises 14.1

**7** For  $f(x, y) = e^{x^2 - y}$ , find the domain and range of  $f$  and the value of  $f(2, 4)$ .

**9** For  $f(x, y, z) = x^2 \ln(x - y + z)$ , find the domain and range of  $f$  and the value of  $f(3, 6, 4)$ .

### Exercises 14.2

Investigate limits of the following expressions at any point at which the expressions is not obviously continuous.

$$\frac{x^2}{x^2 + y^2} \quad (7)$$

$$\frac{xy}{\sqrt{x^2 + y^2}} \quad (11)$$

$$e^{-xy} \sin \pi z / 2 \quad (17)$$

$$\frac{x^2 y^3}{2x^2 + y^2} \quad (35)$$