

## SUMMARY OF THE METHOD OF FROBENIUS

Consider the linear, homogeneous, second order equation:

$$y'' + p(x)y' + q(x)y = 0. \quad (1)$$

Suppose that  $x = 0$  a *regular singular point*:

$$xp(x) = \sum_{n=0}^{\infty} p_n x^n, \quad |x| < R_1, \quad x^2q(x) = \sum_{n=0}^{\infty} q_n x^n, \quad |x| < R_2, \quad R_1, R_2 > 0.$$

Define  $\gamma(r) = r(r - 1) + p_0r + q_0$ ; the *indicial equation* is

$$\gamma(r) = 0, \quad \text{roots } r_1, r_2.$$

**Case (i).**  $r_1$  and  $r_2$  are distinct and do not differ by an integer. There are two linearly independent solutions:

$$y_1(x) = x^{r_1} \sum_{n=0}^{\infty} a_n x^n, \quad y_2(x) = x^{r_2} \sum_{n=0}^{\infty} b_n x^n, \quad a_0 = b_0 = 1. \quad (2)$$

**Case (ii).**  $r_1 = r_2$ . There is one solution  $y_1(x)$  of the form given in (2), and a second solution with the form

$$y_2(x) = y_1(x)(\ln x) + x^{r_1} \sum_{n=1}^{\infty} b_n x^n. \quad (3)$$

**Case (iii).**  $r_1 = r_2 + m$ ,  $m$  a positive integer. There is one solution  $y_1(x)$  as in (2), and a second solution with the form

$$y_2(x) = Cy_1(x)(\ln x) + x^{r_2} \sum_{n=0}^{\infty} b_n x^n, \quad b_0 = 1. \quad (4)$$

The constant  $C$  may or may not be zero. One may assume that  $b_m = 0$ ; see below.

## FURTHER COMMENTS

**1. Normalization.** In these formulas we have “normalized” the solutions by choosing  $a_0$  and  $b_0$  to have value 1. We could just as well have said only that they were nonzero, but it is convenient to have the solutions  $y_1(x)$  and  $y_2(x)$  completely defined.

**2. Radius of convergence.** All the power series in (2)–(4) are guaranteed to have radius of convergence at least as big as the smaller of  $R_1$  and  $R_2$ .

**3. Solution procedure, Case (i).** The coefficients  $a_n$  of the solution  $y_1(x)$  are determined by substituting the given expression (2) for  $y_1(x)$  into (1) and then solving successive equations for  $a_1, a_2, \dots$ . These have the form (before we set  $a_0 = 1$ )

$$\gamma(n + r_1)a_n = \text{a linear combination of } a_0, a_1, \dots, a_{n-1}. \quad (5)$$

The coefficients  $b_n$  of the second solution  $y_2(x)$  in Case (i) are found similarly.

**4. Solution procedure, Cases (ii) and (iii).** In these cases one first finds  $y_1(x)$ . The solution  $y_2(x)$  of (3) or (4) can be written as  $y_2(x) = Cy_1(x)(\ln x) + u(x)$ , where  $C = 1$  in Case (ii) and  $C$  is to be determined in Case (iii), and in each case  $u$  is given by a series. Substituting this form into (1) one finds that  $u(x)$  must satisfy the equation

$$u'' + p(x)u' + q(x)u = \frac{C}{x^2} [y_1(x) - xp(x)y_1(x) - 2xy_1'(x)]. \quad (6)$$

One then substitutes the form of the series for  $u(x)$ , as given in (3) or (4), into (6) and solves for  $b_1, b_2, \dots$  and, in Case (iii), for  $C$ . The general structure of the equations will be similar to (5):

$$\gamma(n + r_2)b_n = \text{a linear combination of } C \text{ and } b_1, b_2, \dots, b_{n-1}. \quad (7)$$

Recall that  $C = 1$  in Case (ii). In Case (iii) the constant  $C$  first appears on the right hand side of (7) when  $m = n$ ; then  $\gamma(m + r_2) = \gamma(r_1) = 0$  so that the left hand side vanishes (and  $b_m$  is not determined). Then  $C$  must be chosen to make the right hand side vanish also.

**5. Additional free constants.** Notice that there is no  $b_0$  coefficient in (3). One could include a  $b_0$  term in the solution, but the value of  $b_0$  would not be determined by the equations;  $b_0$  could be chosen freely. Choosing a nonzero value for the  $b_0$ , however, would amount to adding a multiple of  $y_1(x)$  to the solution  $y_2(x)$  as given in (3).

The situation for Case (iii) is similar. The coefficient  $b_m$  in (4) will not be determined during the solution process, and it is simplest to choose  $b_m = 0$ . Choosing a nonzero value for  $b_m$  again amounts to adding a multiple of  $y_1(x)$  to the solution.

**6. An ordinary point.** An ordinary point of a differential equation may be considered, in some sense, as a special case of a regular singular point. If  $x = 0$  is an ordinary point of (1) then the above analysis applies; one finds that  $\gamma(r) = r(r - 1)$  and hence that  $r_1 = 1$  and  $r_0 = 0$ : we are in Case (iii). However, we already know that in this case there are two linearly independent solutions, as power series in  $x$ , which do not contain  $\ln x$ ; this means that necessarily  $C = 0$ .