THE α -INVARIANTS ON TORIC FANO MANIFOLDS

Jian Song

Department of Mathematics
Columbia University, New York, NY 10027

§1. Introduction

The global holomorphic invariant $\alpha_G(M)$ introduced by Tian[14], Tian and Yau[13] is closely related to the existence of Kähler-Einstein metrics. In his solution to the Calabi conjecture, Yau[19] proved the existence of a Kähler-Einstein metric on compact Kähler manifolds with nonpositive first Chern class. Kähler-Einstein metrics do not always exist in the case when the first Chern class is positive, for there exist known obstructions such as the Futaki invariant. For a compact Kähler manifold M with positive first Chern class, Tian[14] proved that M admits a Kähler-Einstein metric if $\alpha_G(M) > \frac{n}{n+1}$, where $n = \dim M$. In the case of compact complex surfaces, he proved that any compact complex surface with positive first Chern class admits a Kähler-Einstein metric except $CP^2 \# 1\overline{CP^2}$ and $CP^2 \# 2\overline{CP^2}$ [16].

There have been many nice results on the classification of toric Fano manifolds. Mabuchi discovered that if a toric Fano manifold is Kähler-Einstein then the barycenter of the polyhedron defined by its anticanonical divisor is at the origin. V. Batyrev and E. Selivanova [2] estimate the lower bound of α -invariant of symmetric toric Fano manifolds which is sufficient to show the existence of Kähler-Einstein metric.

In this paper, we apply the Tian-Yau-Zelditch expansion of the Bergman kernel on polarized Kähler metrics to approximate plurisubharmonic functions and obtain a formula to calculate the α_G -invariants of all toric Fano manifolds precisely. This gives a generalization of the result by V. Batyrev and E. Selivanova[2] and also this formula confirms the earlier result [12] on the estimates of α invariants on $CP^2 \# 1\overline{CP^2}$ and $CP^2 \# 2\overline{CP^2}$.

Our main theorems are

Theorem 1.1 *If X is a toric Fano manifold of complex dimension n then*

(a) $\alpha_G(X) = 1$ *if X is symmetric, otherwise*

(b) $\alpha_G(X) = \frac{\min_{0 \neq v \in S} \frac{|w_v|}{|v|}}{1 + \min_{0 \neq v \in S} \frac{|w_v|}{|v|}} \leq \frac{1}{2}$.

Corollary 1.1 *If X is a toric Fano manifold, then X is symmetric if and only if $\alpha_G(X) = 1$.*

Theorem 1.2 *If X is a toric Fano manifold, then $\{\alpha_{m,G}(X)\}_{m \geq 1}$ is stationary. More precisely, $\alpha_{m,G}(X) = \alpha_G(X)$ if $m \geq m_0$, where m_0 is the least positive integer such that $m_0 v$ is an integral point and v is the minimizer of $\min_{0 \neq v \in S} \frac{|w_v|}{|v|}$.*

Acknowledgements. The author deeply thanks his advisor, Professor D.H. Phong for his constant encouragement and help. He also thanks Professor Jacob Sturm and Professor Zhiqin Lu for their suggestion on this work. This paper is part of the author's future Ph.D. thesis in Math Department of Columbia University.

§2. Notations

Let us first recall the definition of an invariant $\alpha_G(X)$ introduced by Tian. Let X be an n -dimensional compact complex manifold with positive first Chern class $c_1(X)$ and G a compact subgroup of $Aut(X)$. Choose a G -invariant Kähler metric $g = g_{i\bar{j}}$ on X such that $\omega_g = \frac{\sqrt{-1}}{2\pi} \sum g_{i\bar{j}} dz_i \wedge d\bar{z}_j \in c_1(X)$.

Definition Let $P_G(X, g)$ be the set of all C^2 -smooth G -invariant real-valued functions φ such that $\sup_X \varphi = 0$ and $\omega_g + \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial}\varphi > 0$. The $\alpha_G(X)$ invariant is defined as supremum of all $\lambda > 0$ such that

$$\int_X e^{-\alpha\varphi} \omega^n \leq C(\alpha)$$

for all $\varphi \in P_G(X, g)$, where $C(\alpha)$ is a positive constant depending only on α, g and X .

Let N be a lattice of rank n , $M = Hom(N, \mathbb{Z})$ the dual lattice. $M_R = M \otimes_{\mathbb{Z}} R$, $N_R = N \otimes_{\mathbb{Z}} R$. Let $X = X_{\Sigma}$ be a smooth projective toric n -fold defined by a complete fan Δ of regular cones $\Delta \subset M_R$ and denote $\Delta(i)$ the i -dimensional cone of Δ . We put $T = C^* = \{(t_1, t_2, \dots, t_n) | t_i \in C^*\}$. For $a \in M$ and $b \in N$, we define $\langle a, b \rangle \in \mathbb{Z}$, $\chi^a \in Hom_{alggp}(T, C^*)$ by

$$\begin{aligned} \langle a, b \rangle &= \sum_{i=1}^n a_i b_i, \\ \chi^a((t_1, \dots, t_n)) &= t_1^{a_1} t_2^{a_2} \dots t_n^{a_n}. \end{aligned}$$

For each $\rho \in \Delta(1)$, let b_ρ denote the unique fundamental generator of ρ . We now consider the divisor $K = -\sum_{\rho \in \Delta(1)} D(\rho)$ on $X = X_\Delta$. The following theorem is due to Demazure[4].

Theorem 2.3 *K is a canonical divisor of X_Δ , and the following are equivalent:*

- (a) X_Δ is a toric fano manifold.
- (b) $-K$ is ample.
- (c) $-K$ is very ample.
- (d) $\Sigma_{-K} = \{ a \in M_R \mid \langle a, b_\rho \rangle \leq 1 \text{ for all } \rho \in \Delta(1) \}$ is an n -dimensional compact convex polyhedron whose vertices are exactly $\{a_\tau \mid \tau \in \Delta(n)\}$, where each a_τ denotes the unique element of M such that $\langle a_\tau, b \rangle = 1$ for all fundamental generators b of τ .

The maximal torus $T \subset \text{Aut}(X)$ acting on X has an open dense orbit $U \subset X$, so the normalizer $N(T) \subset \text{Aut}(X)$ of T has a natural action on U . Let $W(X) = N(T)/T$ and we identify the maximal torus $T \subset \text{Aut}(X)$ with an open dense orbit U in X by choosing an arbitrary point $x_0 \in U$, then we have the following splitting short exact sequence

$$1 \rightarrow T \rightarrow N(T) \rightarrow W(X) \rightarrow 1,$$

i.e., an embedding $W(X) \hookrightarrow N(T)$. Denote by $K(T) = (S^1)^n$ the maximal compact subgroup in T . We choose G to be the maximal compact subgroup in $N(T)$ generated by $W(X)$ and $K(T)$, so that we have the short exact sequence

$$1 \rightarrow K(T) \rightarrow G \rightarrow W(X) \rightarrow 1.$$

Proposition 2.1 *Let $X=X_\Delta$ be a smooth projective toric n -fold defined by a complete regular polyhedral fan Δ . Then the group $W(X)$ is isomorphic to the finite group of all symmetries of Δ , i.e., $W(X)$ is isomorphic to a subgroup of $GL(N) (\simeq GL(n, \mathbb{Z}))$ consisting of all elements $\gamma \in GL(N)$ such that $\gamma(\Delta) = \Delta$.*

Remark: $W(X)$ is as well isomorphic to a subgroup of $GL(M) (\simeq GL(n, \mathbb{Z}))$ consisting of all elements $\gamma \in GL(M)$ such that $\gamma(\Sigma) = \Sigma$.

Definition A toric n -fold X is symmetric if the trivial character is the only $W(X)$ -invariant algebraic character of T , i.e. $N^{W(X)} = \{\chi \in N \mid g\chi = \chi \text{ for all } g \in W(X)\} = \{0\}$.

Definition Let $S = \{v \in \partial\Sigma \mid gv = v \text{ for all } g \in W(X)\}$ be the stable points of $W(X)$ on the boundary of Σ . If $S \neq \{0\}$ then for any $0 \neq v \in S$, we define w_v related with v by $w_v = \partial\Sigma \cap \{-tv \mid t \geq 0\}$.

Remark: It's easy to see X is symmetric if and only if $S = 0$.

§3. Holomorphic approximation of PSH

In this section, we will employ the technique in [15, 20] to obtain the approximation of plurisubharmonic functions by logarithms of holomorphic sections of line bundles. The Tian-Yau-Zelditch asymptotic expansion of the potential of the Bergman metric is given by the following theorem[20].

Theorem 3.4 *Let M be a compact complex manifold of dimension n and let $(L, h) \rightarrow M$ be a positive Hermitian holomorphic line bundle. Let g be the Kähler metric on M corresponding to the Kähler form $\omega_g = \text{Ric}(h)$. For each $m \in \mathbb{N}$, h induces a Hermitian metric h_m on L^m . Let $\{S_0^m, S_1^m, \dots, S_{d_m-1}^m\}$ be any orthonormal basis of $H^0(M, L^m)$, $d_m = \dim H^0(M, L^m)$, with respect to the inner product:*

$$(S_1, S_2)_{h_m} = \int_M h_m(S_1(x), S_2(x)) dV_g,$$

where $dV_g = \frac{1}{n!} \omega_g^n$ is the volume form of g . Then there is a complete asymptotic expansion:

$$\sum_{i=0}^{d_m-1} \|S_i^m(x)\|_{h_m}^2 \sim a_0(x)m^n + a_1(x)m^{n-1} + a_2(x)m^{n-2} + \dots$$

for some smooth coefficients $a_j(x)$ with $a_0 = 1$. More precisely, for any k :

$$\left\| \sum_{i=0}^{d_m-1} \|S_i^m(x)\|_{h_m}^2 - \sum_{j < R} a_j(x)m^{n-j} \right\|_{C^k} \leq C_{R,k} m^{n-R}$$

where $C_{R,k}$ depends on R, k and the manifold M .

Let

$$\begin{aligned} \tilde{\omega}_g &= \omega_g + \sqrt{-1} \partial \bar{\partial} \phi > 0 \\ \tilde{h} &= h e^{-\phi} \end{aligned}$$

Let $\tilde{h}_m$ be the induced Hermitian metric of $\tilde{h}$ on L^m , $\{\tilde{S}_0^m, \tilde{S}_1^m, \dots, \tilde{S}_{d_m-1}^m\}$ be any orthonormal basis of $H^0(M, L^m)$, where $d_m = \dim H^0(M, L^m)$, with respect to the inner product

$$(S_1, S_2)_{\tilde{h}_m} = \int_M \tilde{h}_m(S_1(x), S_2(x)) dV_{\tilde{g}}.$$

By Theorem 3.4, we have

$$\sum_{i=0}^{d_m-1} \|\tilde{S}_i^m(x)\|_{h_m}^2 = \left(\sum_{i=0}^{d_m-1} \|\tilde{S}_i^m(x)\|_{h_m}^2 \right) e^{-m\phi}.$$

Thus

$$\phi - \frac{1}{m} \log \left(\sum_{i=0}^{d_m-1} \|\tilde{S}_i^m(x)\|_{h_m}^2 \right) = -\frac{1}{m} \log \left(\sum_{i=0}^{d_m-1} \|\tilde{S}_i^m(x)\|_{h_m}^2 \right)$$

As $m \rightarrow +\infty$, we obtain for any positive integer R

$$\begin{aligned} & \frac{1}{m} \log \left(\sum_{j < R} \tilde{a}_j(x) m^{n-j} \right) \\ &= \frac{1}{m} \log m^n \left(\sum_{j < R} \tilde{a}_j(x) m^{-j} \right) \\ &= \frac{n}{m} \log m + \frac{1}{m} \log(1 + O(\frac{1}{m})) \rightarrow 0 \end{aligned}$$

Thus we have the following corollary of the Tian-Yau-Zelditch expansion.

Corollary 3.2

$$\left\| \phi - \frac{1}{m} \log \left(\sum_{i=0}^{d_m-1} \|\tilde{S}_i^m(x)\|_{h_m}^2 \right) \right\|_{C^k} \rightarrow 0, \text{ as } m \rightarrow +\infty.$$

In other words, any plurisubharmonic function can be approximated by the logarithms of holomorphic sections of L^m .

§4. Proof of The Main Theorem

Suppose X_Δ is Fano, then one obtains a convex $W(X)$ -invariant polyhedron Σ in M_R defined by $\Sigma = \{ a \in M_R \mid \langle a, b_\rho \rangle \leq 1, \text{ for all } \rho \in \Delta(1) \}$ where b_ρ is the fundamental generator of ρ . Let $L(\Sigma) = \{v_0, v_1, \dots, v_k\} = M \cap \Sigma$. Then $v_0, v_1, \dots, v_k$ determine algebraic characters $\chi_i : T \rightarrow C^*$ of T ($i=0, 1, \dots, k$). Moreover, we have

$$|\chi_i(x)|^2 = e^{\langle v_i, y \rangle}, i = 0, \dots, k,$$

where y is the image of x under the canonical projection $\pi : T \rightarrow M_R$. Let us define $u : U \rightarrow R$ as follows:

$$u = \log \left(\sum_{i=0}^k |\chi_i(x)|^2 \right), x \in U \simeq T.$$

Since u is $K(T)$ -invariant, u descends to a function $\tilde{u} : M_R \rightarrow R$ defined as

$$\tilde{u} = \log\left(\sum_{i=0}^k e^{\langle v_i, y \rangle}\right), y \in M_R.$$

Consider the G -invariant hermitian metric $g = g_{i\bar{j}}$ on X such that the restriction of the corresponding to g differential 2-form on U is defined by

$$\omega_g = \partial\bar{\partial}u.$$

The metric g is exactly the pull-back of Fubini-Study metric from P^m with respect to the anticanonical embedding $X \hookrightarrow P^m$ defined by the algebraic characters $\chi_0, \chi_1, \dots, \chi_k$.

Let $\Sigma^{(m)} = \{a \in M_R \mid \langle a, b_\rho \rangle \leq m \text{ and } L(\Sigma^{(m)}) = \{v_0, \dots, v_{k_m}\} = M \cap \Sigma^{(m)}, \text{ where } k_m + 1 = \dim H^0(X, O((-K)^m)) \text{ and } \chi^\mu : T \rightarrow C^* \text{ defined by } |\chi^\mu(x)|^2 = e^{\langle \mu, y \rangle}\}$. We have the following lemma (see [7] p66)

Lemma 4.1 $H^0(X, O((-K)^m)) = \oplus_{\mu \in L(\Sigma^{(m)})} C \cdot \chi^\mu$.

Proposition 4.2 $\{\chi^\mu\}_{\mu \in L(\Sigma^{(m)})}$ is an orthogonal basis of $H^0(X, O((-K)^m))$ with respect to the inner product $\langle, \rangle_{h^m}$, where $h^m = \frac{1}{(\sum_{i=0}^k |\chi_i(x)|^2)^m}$.

Proof

$$\begin{aligned} & \int_X \langle \chi^\mu, \chi^\nu \rangle_{h^m} \omega^n \\ &= \int_T \frac{(z_1^{\mu_1} \dots z_n^{\mu_n})(\bar{z}_1^{\nu_1} \dots \bar{z}_n^{\nu_n})}{(\sum_{i=0}^k |z^{v_i}|^m)} \omega^n \\ &= \int_T \frac{|z_1|^{\mu_1 + \nu_1} \dots |z_n|^{\mu_n + \nu_n} e^{i(\mu_1 - \nu_1)\theta_1} \dots e^{i(\mu_n - \nu_n)\theta_n}}{(\sum_{i=0}^k |z^{v_i}|^m)} \omega^n. \end{aligned}$$

which is 0 if $\mu \neq \nu$.

For any $\varphi \in P_G(X, \omega)$, by Corollary 3.3

$$\varphi(x) = \lim_{m \rightarrow \infty} \frac{1}{m} \log \frac{\sum_{\mu \in L(\Sigma^{(m)})} a_\mu^{(m)} |\chi^\mu(x)|^2}{(\sum_{i=0}^k |\chi_i(x)|^2)^k}$$

Lemma 4.2 There exists $\epsilon > 0$ such that for $\varphi \in P_G(X, \omega)$ and $\tilde{m} > 0$ there exist $m > \tilde{m}$ and $\mu \in L(\Sigma^{(m)})$ with $(a_\mu^{(m)})^{\frac{1}{m}} > \epsilon$.

Proof Otherwise, for any $\epsilon > 0$ there exists φ_ϵ and $\tilde{m}$ such that for any $m > \tilde{m}$ and $\mu \in L(\Sigma^{(m)})$ we have $(a_\mu^{(m)})^{\frac{1}{m}} < \epsilon$. By choosing m large enough we have

$$\begin{aligned}
\varphi_\epsilon(x) &\leq \frac{1}{m} \log \frac{\sum_{\mu \in L(\Sigma^{(m)})} |\chi^\mu(x)|^2}{(\sum_{i=0}^k |\chi_i(x)|^2)^m} + \log \epsilon \\
&= \frac{1}{m} \log \left(\sum_{\mu \in L(\Sigma^{(m)})} \frac{|\chi^\mu(x)|^2}{(\sum_{i=0}^k |\chi_i(x)|^2)^m} \right) + \log \epsilon \\
&\leq \frac{1}{m} \log \left(\sum_{\mu \in L(\Sigma^{(m)})} 1 \right) + \log \epsilon \\
&\leq \text{Const} + \log \epsilon.
\end{aligned}$$

Since ϵ can be chosen arbitrarily small, the above inequality implies that $\varphi_\epsilon \rightarrow -\infty$ uniformly as ϵ goes to 0, which contradicts the fact that $\sup_X \varphi = 0$.

For any $\varphi \in P_G(X, \omega)$, by Lemma 4.1 we have

$$\begin{aligned}
\varphi(x) &= \lim_{m \rightarrow \infty} \frac{1}{m} \log \frac{\sum_{\mu \in L(\Sigma^{(m)})} a_\mu^{(m)} |\chi^\mu(x)|^2}{(\sum_{i=0}^k |\chi_i(x)|^2)^m} \\
&\geq \frac{1}{m} \log \frac{\sum_{g \in W(X)} |\chi^{g\mu}(x)|^2}{(\sum_{i=0}^k |\chi_i(x)|^2)^m} - C_1 \\
&\geq \log \frac{|\chi^{\sum_{g \in W(X)} g\mu}(x)|^{\frac{2}{m|W(X)|}}}{(\sum_{i=0}^k |\chi_i(x)|^2)} - C_1 \\
&\geq \log \frac{|\chi(x)|^{\frac{2 \sum_{g \in W(X)} g\mu}{m|W(X)|}}}{(\sum_{i=0}^k |\chi_i(x)|^2)} - C_1
\end{aligned}$$

Put $v = \frac{\sum_{g \in W(X)} g\mu}{|W(X)|K}$, then we have $\tilde{\varphi}(y) \geq \log \frac{e^{\langle v, y \rangle}}{\sum_{i=0}^m e^{\langle v_i, y \rangle}}$

Put $y_i = \log |t_i|^2$ $t_i = e^{\frac{y_i}{2} + \sqrt{-1}\theta_i}$, then

$$\begin{aligned}
\frac{dt_i}{t_i} &= \frac{1}{2} dy_i + \sqrt{-1} d\theta_i \\
\frac{d\bar{t}_i}{\bar{t}_i} &= \frac{1}{2} dy_i + \sqrt{-1} d\theta_i \\
\frac{dt_i \wedge d\bar{t}_i}{|t_i|^2} &= -\sqrt{-1} dy_i \wedge d\theta_i \\
\partial \bar{\partial} u &= \sum_{i,j} \frac{\partial^2 u}{\partial y_i \partial y_j} \frac{dt_i \wedge d\bar{t}_j}{t_i \bar{t}_j} \\
(\partial \bar{\partial} u)^n &= \det \left(\frac{\partial^2 u}{\partial y_i \partial y_j} \right) dy_1 \wedge \dots \wedge dy_n \wedge d\theta_1 \wedge \dots \wedge d\theta_n
\end{aligned}$$

Lemma 4.3 Let $\tilde{F} = e^{\tilde{u}} \det \frac{\partial^2 \tilde{u}}{\partial y_i \partial y_j}$, then $0 < c \leq \tilde{F} \leq C$.

Proof $e^{-u} \frac{dt_1 \wedge d\bar{t}_1 \wedge \dots \wedge dt_n \wedge d\bar{t}_n}{|t_1|^2 \dots |t_n|^2} = e^{-\tilde{u}} dy_1 \wedge \dots \wedge dy_n \wedge d\theta_1 \wedge \dots \wedge d\theta_n$ can be extended to a non-vanishing volume form on X . Also

$$\begin{aligned} (\partial \bar{\partial} u)^n &= \det \frac{\partial^2 u}{\partial t_i \partial \bar{t}_j} dt_1 \wedge d\bar{t}_1 \wedge \dots \wedge dt_n \wedge d\bar{t}_n \\ &= \det \frac{\partial^2 \tilde{u}}{\partial y_i \partial y_j} dy_1 \wedge \dots \wedge dy_n \wedge d\theta_1 \wedge \dots \wedge d\theta_n \end{aligned}$$

is a non-vanishing volume form, so the quotient of these two volume form must be positive and bounded. which proves the lemma.

Now we can prove the Theorem 1.1.

$$\begin{aligned} \int_X e^{-\alpha \varphi} \omega^n &= \int_X e^{-\alpha \varphi} (\partial \bar{\partial} u)^n \\ &= \int_{R^n} e^{-\alpha \tilde{\varphi}} \det \frac{\partial^2 \tilde{u}}{\partial y_i \partial y_j} dy_1 \dots dy_n \\ &\leq \int_{R^n} e^{-\alpha \tilde{\varphi} - \tilde{u}} dy_1 \dots dy_n \\ &\leq \int_{R^n} e^{-\alpha \log \sum e^{<v_i, y>} - \log(\sum e^{<v_i, y>})} dy_1 \dots dy_n \\ &= \int_{R^n} \frac{e^{-\alpha <v, y>}}{(\sum e^{<v_i, y>})^{1-\alpha}} dy_1 \dots dy_n \end{aligned}$$

If the stable points $S = \{0\}$, then X is symmetric so that $v = \frac{\sum_{g \in W(X)} g\mu}{m|W(X)|} = 0$ for all $\mu \in L(\Sigma^{(m)})$. Therefore for all $\alpha < 1$ the integral

$$\int_X e^{-\alpha \varphi} \omega^n = \int_{R^n} \frac{1}{(\sum_{i=0}^k e^{<v_i, y>})^{1-\alpha}} dy_1 \dots dy_n$$

is finite since every n -dimensional cone $\sigma_j \in \Delta (j = 1, \dots, l)$ is generated by a basis of the lattice N and the fact that $N_R = \sigma_1 \cup \dots \cup \sigma_l$. This implies $\alpha_G(X) \geq 1$ so that by Tian's theorem[14] X admits Kahler-Einstein metric. This is a result by V. Batyrev and E.N. Selivanova[2].

If $S \neq \{0\}$ then for any $0 \neq v \in S$, we have $w_v \in \partial \Sigma$ related with v satisfying

$$< w_v, v > = -|w_v||v|.$$

The integral

$$\int_{R^n} \frac{e^{-\alpha \langle v, y \rangle}}{(\sum e^{\langle v_i, y \rangle})^{1-\alpha}} dy_1 \dots dy_n = \int_{R^n} \left(\frac{e^{-\frac{\alpha}{1-\alpha} \langle v, y \rangle}}{\sum e^{\langle v_i, y \rangle}} \right)^{1-\alpha} dy_1 \dots dy_n$$

is finite if

$$-\frac{\alpha}{1-\alpha}v \in \text{int}(\Sigma).$$

i.e.

$$\langle -\frac{\alpha}{1-\alpha}v, w_v \rangle = \frac{\alpha}{1-\alpha}|v||w_v| \leq |w_v|^2$$

Then for all $\alpha < \frac{\min_{0 \neq v \in S} \frac{|w_v|}{|v|}}{1 + \min_{0 \neq v \in S} \frac{|w_v|}{|v|}}$ the integral $\int_X e^{-\alpha \varphi} \omega^n$ is finite. Therefore

$$\alpha_G(X) \geq \frac{\min_{0 \neq v \in S} \frac{|w_v|}{|v|}}{1 + \min_{0 \neq v \in S} \frac{|w_v|}{|v|}}.$$

In order to estimate the upper bound of $\alpha_G(X)$ we will construct a sequence of PSH functions. Suppose $S \neq \{0\}$, then for all α with $1 > \alpha > \frac{\min_{0 \neq v \in S} \frac{|w_v|}{|v|}}{1 + \min_{0 \neq v \in S} \frac{|w_v|}{|v|}}$ we choose $\tilde{\varphi}_\epsilon = \log(\frac{e^{\langle \tilde{v}, y \rangle + \epsilon}}{\sum e^{\langle v_i, y \rangle}})$ which is increasing and uniformly bounded from above where $\min_{0 \neq v \in S} \frac{|w_v|}{|v|}$ is achieved at $\tilde{v} \in S$. Then by Fatou lemma we have

$$\lim_{\epsilon \rightarrow 0} \int_X e^{-\alpha \varphi_\epsilon} \omega^n = \int_X e^{-\alpha \varphi_0} \omega^n = \infty$$

which implies $\alpha_G(X) \leq \frac{\min_{0 \neq v \in S} \frac{|w_v|}{|v|}}{1 + \min_{0 \neq v \in S} \frac{|w_v|}{|v|}}$. Combined the above estimates together, we have proved the Theorem 1.1.

Also it's obvious to see that $\min_{0 \neq v \in S} \frac{|w_v|}{|v|} \leq 1$ for non-symmetric toric Fano manifold X thus $\alpha_G(X) \leq \frac{1}{2}$. This shows that there doesn't exist any non-symmetric toric Fano manifold such that its α_G -invariant is greater than $\frac{n}{n+1}$ which is a sufficient condition for the existence of Kähler-Einstein metrics.

Now we prove Theorem 1.2 which is a direct corollary of the proof of Theorem 1.1. Define $P_{m,G}(X) = \{\varphi \in C^\infty(X, \mathbb{R}) \mid \sup_X \varphi = 0, \varphi \text{ is } G\text{-invariant and there exists a basis } \{S_i^m\}_{1 \leq i \leq N_m} \text{ of } H^0(X, K_X^{-m}) \text{ such that } \omega_g + \partial \bar{\partial} \varphi = \frac{1}{m} \partial \bar{\partial} \log(\sum_{i=0}^{N_m} |S_i^m|^2)\}$, where $N_m + 1 = \dim H^0(X, K_X^{-m})$ and m is large.

We also define for m large, $\alpha_{m,G}(X) = \sup\{\alpha \mid \text{there exists } C > 0 \text{ such that for all } \varphi \in P_{m,G}(X), \int_X e^{-\alpha \varphi} dV \leq \infty\}$.

It's easy to see $\alpha_{m,G}(X)$ is decreasing as m goes to the infinity. By the argument to give the upper bound for the α_G -invariant, we can directly have the following corollary which answers the question proposed by Tian[15] in the special case of toric Fano manifolds.

Corollary 4.3 *If X is a toric Fano manifold, then $\{\alpha_{m,G}(X)\}_m$ is decreasing and stationary. More precisely, $\alpha_{m,G}(X) = \alpha_G(X)$ if $m \geq m_0$, where m_0 is the least positive integer such that $m_0 v \in M$ and v is the minimizer of $\min_{0 \neq v \in S} \frac{|w_v|}{|v|}$.*

§5. Multiplier ideal sheaf

In this section we relate the α -invariant on toric Fano manifolds with the method of the multiplier ideal sheaf employed by Nadel[9]. Here we follow the lines in [3].

Theorem 5.5 (Nadel) *Let X be a Fano manifold of dimension n and G be a compact subgroup of the group of complex automorphisms of X . Then X admits a G -invariant Kähler-Einstein metric, unless K_X^{-1} possesses a G -invariant singular hermitian metric $h = h_0 e^{-\varphi}$ (h_0 is a smooth G -invariant metric and φ is a G -invariant function in $L^1_{loc}(X)$), such that the following properties occur.*

1. h has a semipositive curvature current

$$\Theta_h = -\frac{i}{2\pi} \partial \bar{\partial} \log h = \Theta_{h_0} + \frac{i}{2\pi} \partial \bar{\partial} \varphi \leq 0.$$

2. For every $\gamma \in (\frac{n}{n+1}, 1)$, the multiplier ideal sheaf $\mathcal{J}(\gamma\varphi)$ is nontrivial, (i.e. $0 \neq \mathcal{J}(\gamma\varphi) \neq \mathcal{O}_X$).

Theorem 5.6 (Nadel) *Let (X, ω) be a Kähler manifold and let L be a holomorphic line bundle over X equipped with a singular hermitian metric h of weight ϕ with respect to a smooth metric h_0 (i.e. $h = h_0 e^{-\phi}$). Assume that the curvature form $\Theta_h(L)$ is positive definite in the sense of currents, i.e. $\Theta_h(L) \geq \epsilon \omega$ for some $\epsilon > 0$. Then we have $H^q(X, K_X \otimes L \otimes \mathcal{J}(\phi)) = 0$ for all $q \geq 1$*

Corollary 5.4 (Nadel) *Let X , G , h and φ be in theorem 4.1. then for all $\gamma \in (\frac{n}{n+1}, 1)$,*

1. *The multiplier ideal sheaf $\mathcal{J}(\gamma\varphi)$ satisfies $H^q(X, \mathcal{J}(\gamma\varphi)) = 0$ for all $q \geq 1$.*
2. *The associated subscheme V_γ of structure sheaf $\mathcal{O}_{V_\gamma} = \mathcal{O}_X / \mathcal{J}(\gamma\varphi)$ is nonempty, distinct from X , G -invariant and satisfies $H^q(V_\gamma, \mathcal{O}_{V_\gamma}) = C$ for $q=0$ and vanishes for $q \geq 1$.*

In order to construct Kahler-Einstein metrics it's sufficient to rule out the existence of any G -invariant subscheme with the property (2) in the corollary.

In the case of toric Fano manifolds, we have the following theorem.

Theorem 5.7 *Let X be a toric Fano manifold. If X is not symmetric then there always exists a G -invariant subscheme with the property (2) in the corollary.*

Proof If X is not symmetric then $\alpha_G(X) \leq \frac{1}{2}$ and we can construct G -invariant $\varphi \in L_{loc}^1(X)$ such that for all $\gamma \in (\alpha_G(X), 1)$

$$\int_X e^{-\gamma\varphi} \omega^n = +\infty$$

therefore $J(\gamma\varphi)$ is nontrivial and there exist subschemes V_γ satisfying property (2) of the corollary.

§6. Examples

In this sections we will calculate the α invariants for 2-dimensional toric Fano manifolds.

Here (1) (2) (3) (4) are corresponding to CP^2 and CP^2 blow-up at 1, 2 and 3 points and (5) (6) (7) (8) are the corresponding polyhedrons.

CP^2 and CP^2 blow-up at 3 points are symmetric thus the their α_G -invariants are both equal to 1.

For $CP^2 \# 1\overline{CP^2}$ its stable points of G on the boundary of the polyhedron in (6) is $(\frac{1}{2}, \frac{1}{2})$ and $(-\frac{1}{2}, -\frac{1}{2})$ and $\frac{|(1/2, 1/2)|}{|(-1/2, -1/2)|} = 1$ then it is easy to see $\alpha_G(CP^2 \# 1\overline{CP^2}) = \frac{1}{2}$.

For $CP^2 \# 2\overline{CP^2}$ its stable points of G on the boundary of the polyhedron in (7) is $(\frac{1}{2}, \frac{1}{2})$ and $(-1, -1)$ and $\frac{|(1/2, 1/2)|}{|(-1, -1)|} = \frac{1}{2}$ then it is easy to see $\alpha_G(CP^2 \# 2\overline{CP^2}) = \frac{1}{3}$.

The above calculation confirms our earlier results[12].

References

- [1] Abdesselem, B. A., *Équations de Monge-Ampère d'origine géométrique sur certaines variétés algébriques*, J.Funct. Anal. 149 (1997), 102-134 Math. Soc. (3), 50 (1985), 1-26.
- [2] Batyrev, V. and Selivanova, E., *Einstein-Kähler metrics on symmetric toric Fano manifolds*, J. reine angew. Math. 512 (1999), 225-236.
- [3] Demailly, J.-P. and Kollár, J., *Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds*, Ann. Sci. école Norm. Sup. (4) 34 (2001), no. 4, 525–556.
- [4] Demazure, M., *sous-groupes algébriques de rang maximum du groupe de Cremona*, Ann. Sci. École Norm. Sup. 3 (1970), 507-588.
- [5] Danilov, V.I., *Geometry of toric varieties*, Russ. Math. Surv. 33, No.2 (1978), 97-154.
- [6] Ding, W and Tian, G., *The generalized Moser-Trudinger Inequality*, Proceedings of Nankai International Conference on Nonlinear Analysis, 1993.
- [7] Fulton, W., *Introduction to Toric Varieties*, Ann. Math. Stud. 131, Princeton Univ. Press, 1993.
- [8] Mabuchi, T., *Einstein-Kähler forms, Futaki invariants and convex geometry on toric Fano varieties*, Osaka J. Math. 24 (1987), 705-737.
- [9] Nadel, A.M., *Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature*, Ann. of Math. (2) 132 (1990), no. 3, 549–596.
- [10] Phong, D. H. and Sturm, J., *Algebraic estimates, stability of local zeta functions, and uniform estimates for distribution functions*, Ann. of Math. (2) 152 (2000), no. 1, 277–329.
- [11] Siu, Y.T., *The existence of Kähler-Einstein metrics on manifolds with positive anticanonical line bundle and a suitable finite symmetry group*, Ann. of Math. (2) 127 (1988), no. 3, 585–627.
- [12] Song, J., *The α -Invariant on $CP^2 \# 2\overline{CP^2}$* , preprint.

- [13] Tian, G. and Yau, S.T., *Kähler-Einstein metrics on complex surfaces with $C_1 > 0$* , Comm. Math. Phys. 112 (1987), no.1,175–203.
- [14] Tian, G., *On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$* , Invent. Math. 89 (1987), no. 2, 225–246.
- [15] Tian, G., *On a set of polarized Kähler metrics on algebraic manifolds*, J.Differential Geometry 32(1990)99-130.
- [16] Tian, G., *On Calabi's conjecture for complex surfaces with positive first Chern class*, Invent. Math. 101 (1990), no. 1, 101–172.
- [17] Tian, G., *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. 130 (1997), no. 1, 1–37.
- [18] Tian, G. and Zhu, X.H., *A nonlinear inequality of Moser-Trudinger type*, Calc. Var. Partial Differential Equations 10 (2000), no. 4, 349–354.
- [19] Yau, S.T., *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I*, Comm. Pure Appl. Math. 31 (1978) 339-411.
- [20] Zelditch, S., *Szego Kernels and a Theorem of Tian*, IMRN1998, No.6 317-331.