

A Smooth Converse Lyapunov Theorem for Robust Stability

Yuandan Lin*

Department of Mathematics
Florida Atlantic University
Boca Raton, FL 33431

yuandan@polya.math.fau.edu

Eduardo D. Sontag*

Department of Mathematics
Rutgers University
New Brunswick, NJ 08903

sontag@hilbert.rutgers.edu

Yuan Wang[†]

Department of Mathematics
Florida Atlantic University
Boca Raton, FL 33431

ywang@polya.math.fau.edu

Abstract. This paper presents a Converse Lyapunov Function Theorem motivated by robust control analysis and design. Our result is based upon, but generalizes, various aspects of well-known classical theorems. In a unified and natural manner, it (1) allows arbitrary bounded time-varying parameters in the system description, (2) deals with global asymptotic stability, (3) results in smooth (infinitely differentiable) Lyapunov functions, and (4) applies to stability with respect to not necessarily compact invariant sets.

1. Introduction. This work is motivated by problems of robust nonlinear stabilization. One of our main contributions is to provide a statement and proof of a Converse Lyapunov Function Theorem which is in a form particularly useful for the study of such feedback control analysis and design problems. We provide a single (and natural) unified result that:

1. applies to stability with respect to not necessarily compact invariant sets;
2. deals with global (as opposed to merely local) asymptotic stability;
3. results in smooth (infinitely differentiable) Lyapunov functions;
4. most importantly, applies to stability in the presence of bounded time varying parameters in the system.

(This latter property is sometimes called “total stability” and it is equivalent to the stability of an associated differential inclusion.)

The interest in stability with respect to possibly non-compact sets is motivated by applications to areas such as output-control (one needs to stabilize with respect to the zero set of the output variables) and Luenberger-type observer design (“detectability” corresponds to stability with respect to the diagonal set $\{(x, x)\}$, as a subset of the composite state/observer system). Such applications and others are explored in [16], Chapter 5.

Smooth Lyapunov functions, as opposed to merely continuous or once-differentiable, are required in order to apply “backstepping” techniques in which a feedback law is built by successively taking directional derivatives of feedback laws obtained for a simplified system. (See for instance [9] for more on backstepping design.)

Finally, the effect of parameter uncertainty, and the study of associated Lyapunov functions, are topics of interest in robust control theory. An application of the result proved in this paper to the study of “input to state stability” is provided in [27].

* Supported in part by US Air Force Grant AFOSR-91-0346

[†] Supported in part by NSF Grant DMS-9108250

Keywords: Nonlinear stability, Stability with respect to sets, Lyapunov function techniques, Robust stability.

Running head: Converse Lyapunov Theorem for Robust Stability

AMS(MOS) subject classifications: 93D05, 93D09, 93D20, 34D20

1.1. Organization of Paper. The paper is organized as follows.

The next section provides the basic definitions and the statement of the main result. Actually, two versions are given, one that applies to global asymptotic stability with respect to arbitrary invariant sets, but assuming completeness of the system—that is, global existence of solutions for all inputs—and another version which does not assume completeness but only applies to the special case of compact invariant sets (in particular, to the usual case of global asymptotic stability with respect to equilibria).

Equivalent characterizations of stability by means of decay estimates have proved very useful in control theory—see e.g. [25]—and this is the subject of Section 3. Some technical facts about Lyapunov functions, including a result on the smoothing of such functions around an attracting set, are given in Section 4. After this, Section 5 establishes some basic facts about complete systems needed for the main result.

Section 6 contains the proof of the main result for the general case. Our proof is based upon, and follows to a great extent the outline of, the one given by Wilson in [31], who provided in the late 1960s a converse Lyapunov function theorem for local asymptotic stability with respect to closed sets. There are however some major differences with that work: we want a global rather than a local result, and several technical issues appear in that case; moreover, and most importantly, we have to deal with parameters, which makes the careful analysis of uniform bounds of paramount importance. (In addition, even for the case of no parameters and local stability, several critical steps in the proof are only sketched in [31], especially those concerning Lipschitz properties and smoothness around the attracting set. Later the author of [21] rederived the results, but only for the case when the invariant set is compact. Thus it seems useful to have an expository detailed and self-contained proof in the literature for the more general cases.) A needed technical result on smoothing functions, based closely also on [31], is placed in an Appendix for convenience. Section 7 deals with the compact case, essentially by reparameterization of trajectories.

An example, motivated by related work of Tsinias and Kalouptsidis, is given in Section 8 to show that the analogous theorems are false for unbounded parameters.

Obviously in a topic such as this one, there are many connections to previous work. While it is likely that we have missed many relevant references, we discuss in Section 9 some relationships between our work and other results in the literature. Relations to work using “prolongations” are particularly important, and are the subject of some more detail in Section 10.

2. Definitions and Statements of Main Results. Consider the following system:

$$(1) \quad \dot{x}(t) = f(x(t), d(t)),$$

where for each $t \in \mathbb{R}$, $x(t) \in \mathbb{R}^n$ and $d(t) \in \mathcal{D}$, and where \mathcal{D} is a compact subset of \mathbb{R}^m , for some positive integers n and m . The map $f : \mathbb{R}^n \times \mathcal{D} \rightarrow \mathbb{R}^n$ is assumed to satisfy the following two properties:

- f is continuous.
- f is locally Lipschitz on x uniformly on d , that is, for each compact subset K of \mathbb{R}^n there is some constant c so that $|f(x, \mathbf{d}) - f(z, \mathbf{d})| \leq c|x - z|$ for all $x, z \in K$ and all $\mathbf{d} \in \mathcal{D}$, where $|\cdot|$ denotes the usual Euclidian norm.

Note that these properties are satisfied, for instance, if f extends to a continuously differentiable function on a neighborhood of $\mathbb{R}^n \times \mathcal{D}$.

Let $\mathcal{M}_{\mathcal{D}}$ be the set of all measurable functions from \mathbb{R} to \mathcal{D} . We will call functions $d \in \mathcal{M}_{\mathcal{D}}$ *time varying parameters*. For each $d \in \mathcal{M}_{\mathcal{D}}$, we denote by $x(t, x_0, d)$ (and sometimes simply by $x(t)$ if there is no ambiguity from the context) the solution at time t of (1) with $x(0) = x_0$. This is defined on some maximal interval $(T_{x_0, d}^-, T_{x_0, d}^+)$ with $-\infty \leq T_{x_0, d}^- < 0 < T_{x_0, d}^+ \leq +\infty$.

Sometimes we will need to consider time varying parameters d that are defined only on some interval $I \subseteq \mathbb{R}$ with $0 \in I$. In those cases, by abuse of notation, $x(t, x_0, d)$ will still be used, but only times $t \in I$ will be considered.

The system is said to be *forward complete* if $T_{x_0,d}^+ = +\infty$ for all x_0 all $d \in \mathcal{M}_{\mathcal{D}}$. It is *backward complete* if $T_{x_0,d}^- = -\infty$ for all x_0 all $d \in \mathcal{M}_{\mathcal{D}}$, and it is *complete* if it is both forward and backward complete.

We say that a closed set \mathcal{A} is an *invariant set* for (1) if

$$\forall x_0 \in \mathcal{A}, \forall d \in \mathcal{M}_{\mathcal{D}}, T_{x_0,d}^+ = +\infty \text{ and } x(t, x_0, d) \in \mathcal{A}, \forall t \geq 0.$$

REMARK 2.1. An equivalent formulation of invariance is in terms of the associated differential inclusion

$$(2) \quad \dot{x} \in F(x),$$

where $F(x) = \{f(x, \mathbf{d}), \mathbf{d} \in \mathcal{D}\}$. The set \mathcal{A} is invariant for (1) if and only if it is invariant with respect to (2) (see e.g. [1]). The notions of stability to be considered later can be rephrased in terms of (2) as well. \square

We will use the following notation: for each nonempty subset \mathcal{A} of \mathbb{R}^n , and each $\xi \in \mathbb{R}^n$, we denote

$$|\xi|_{\mathcal{A}} \stackrel{\text{def}}{=} d(\xi, \mathcal{A}) = \inf_{\eta \in \mathcal{A}} d(\xi, \eta),$$

the common point-to-set distance, and $|\xi|_{\{0\}} = |\xi|$ is the usual norm.

Let $\mathcal{A} \subseteq \mathbb{R}^n$ be a closed, invariant set for (1). We emphasize that we do not require \mathcal{A} to be compact. We will assume throughout this work that the following mild property holds:

$$(3) \quad \sup_{\xi \in \mathbb{R}^n} \{|\xi|_{\mathcal{A}}\} = \infty.$$

This is a minor technical assumption, satisfied in all examples of interest, which will greatly simplify our statements and proofs. (Of course, this property holds automatically whenever \mathcal{A} is compact, and in particular in the important special case in which \mathcal{A} reduces to an equilibrium point.)

DEFINITION 2.2. *System (1) is (absolutely) uniformly globally asymptotically stable (UGAS) with respect to the closed invariant set \mathcal{A} if it is forward complete and the following two properties hold:*

1. *Uniform Stability. There exists a \mathcal{K}_{∞} -function $\delta(\cdot)$ such that for any $\varepsilon \geq 0$,*

$$(4) \quad |x(t, x_0, d)|_{\mathcal{A}} \leq \varepsilon \text{ for all } d \in \mathcal{M}_{\mathcal{D}}, \text{ whenever } |x_0|_{\mathcal{A}} \leq \delta(\varepsilon) \text{ and } t \geq 0.$$

2. *Uniform Attraction. For any $r, \varepsilon > 0$, there is a $T > 0$, such that for every $d \in \mathcal{M}_{\mathcal{D}}$,*

$$(5) \quad |x(t, x_0, d)|_{\mathcal{A}} < \varepsilon$$

whenever $|x_0|_{\mathcal{A}} < r$ and $t \geq T$. \square

For the definitions of the standard comparison classes of \mathcal{K}_{∞} and \mathcal{KL} functions, we refer the reader to the appendix.

Observe that when \mathcal{A} is compact the forward completeness assumption is redundant, since in that case property (4) already implies that all solutions are bounded.

In the particular case in which the set \mathcal{D} consists of just one point, the above definition reduces to the standard notion of set asymptotic stability of differential equations. (Note, however, that this definition differs from those in [3], and [31], which are not global.) If, in addition, \mathcal{A} consists of just an equilibrium point x_0 , this is the usual notion of global asymptotic stability for the solution $x(t) \equiv x_0$.

REMARK 2.3. It is an easy exercise to verify that an equivalent definition results if one replaces $\mathcal{M}_{\mathcal{D}}$ by the subset of piecewise constant time varying parameters. \square

REMARK 2.4. Note that the uniform stability condition is equivalent to: there is a \mathcal{K}_{∞} -function φ so that

$$|x(t, x_0, d)|_{\mathcal{A}} \leq \varphi(|x_0|_{\mathcal{A}}), \quad \forall x_0, \forall t \geq 0, \text{ and } \forall d \in \mathcal{M}_{\mathcal{D}}.$$

(Just let $\varphi = \delta^{-1}$.) \square

The following characterization of the UGAS property will be extremely useful.

PROPOSITION 2.5. *The system (1) is UGAS with respect to a closed, invariant set $\mathcal{A} \subseteq \mathbb{R}^n$ if and only if it is forward complete and there exists a \mathcal{KL} -function β such that, given any initial state x_0 , the solution $x(t, x_0, d)$ satisfies*

$$(6) \quad |x(t, x_0, d)|_{\mathcal{A}} \leq \beta(|x_0|_{\mathcal{A}}, t), \quad \text{any } t \geq 0,$$

for any $d \in \mathcal{M}_{\mathcal{D}}$.

Observe that when \mathcal{A} is compact the forward completeness assumption is again redundant, since in that case property (6) implies that solutions are bounded.

Next we introduce Lyapunov functions with respect to sets. For any differentiable function $V : \mathbb{R}^n \rightarrow \mathbb{R}$, we use the standard Lie derivative notation

$$L_{f_{\mathbf{d}}} V(\xi) \stackrel{\text{def}}{=} \frac{\partial V(\xi)}{\partial x} \cdot f_{\mathbf{d}}(\xi),$$

where for each $\mathbf{d} \in \mathcal{D}$, $f_{\mathbf{d}}(\cdot)$ is the vector field defined by $f(\cdot, \mathbf{d})$. By “smooth” we always mean infinitely differentiable.

DEFINITION 2.6. *A Lyapunov function for the system (1) with respect to a nonempty, closed, invariant set $\mathcal{A} \subseteq \mathbb{R}^n$ is a function $V : \mathbb{R}^n \rightarrow \mathbb{R}$ such that V is smooth on $\mathbb{R}^n \setminus \mathcal{A}$ and satisfies*

1. *there exist two \mathcal{K}_{∞} -functions α_1 and α_2 such that for any $\xi \in \mathbb{R}^n$,*

$$(7) \quad \alpha_1(|\xi|_{\mathcal{A}}) \leq V(\xi) \leq \alpha_2(|\xi|_{\mathcal{A}});$$

2. *there exists a continuous, positive definite function α_3 such that for any $\xi \in \mathbb{R}^n \setminus \mathcal{A}$, and any $\mathbf{d} \in \mathcal{D}$,*

$$(8) \quad L_{f_{\mathbf{d}}} V(\xi) \leq -\alpha_3(|\xi|_{\mathcal{A}}).$$

A smooth Lyapunov function is one which is smooth on all of \mathbb{R}^n . □

REMARK 2.7. Continuity of V on $\mathbb{R}^n \setminus \mathcal{A}$ and property 1. in the definition imply:

- V is continuous on all of \mathbb{R}^n ;
- $V(x) = 0 \iff x \in \mathcal{A}$; and
- $V : \mathbb{R}^n \xrightarrow{\text{onto}} \mathbb{R}_{\geq 0}$ (recall the assumption in equation (3)). □

Our main results will be two converse Lyapunov theorems. The first one is for general closed invariant sets and assumes completeness of the system.

THEOREM 2.8. *Assume that the system (1) is complete. Let $\mathcal{A} \subseteq \mathbb{R}^n$ be a nonempty, closed invariant subset for this system. Then, (1) is UGAS with respect to \mathcal{A} if and only if there exists a smooth Lyapunov function V with respect to \mathcal{A} .*

The following result does not assume completeness but instead applies only to compact \mathcal{A} :

THEOREM 2.9. *Let $\mathcal{A} \subseteq \mathbb{R}^n$ be a nonempty, compact invariant subset for the system (1). Then, (1) is UGAS with respect to \mathcal{A} if and only if there exists a smooth Lyapunov function V with respect to \mathcal{A} .*

3. Some Preliminaries about UGAS. It will be useful to have a restatement of the second condition in the definition of UGAS stated in terms of uniform attraction times:

LEMMA 3.1. *The uniform attraction property defined in Definition 2.2 is equivalent to the following: There exists a family of mappings $\{T_r\}_{r>0}$ with*

- *for each fixed $r > 0$, $T_r : \mathbb{R}_{>0} \xrightarrow{\text{onto}} \mathbb{R}_{>0}$ is continuous and is strictly decreasing;*
- *for each fixed $\varepsilon > 0$, $T_r(\varepsilon)$ is (strictly) increasing as r increases and $\lim_{r \rightarrow \infty} T_r(\varepsilon) = \infty$;*

such that, for each $d \in \mathcal{M}_{\mathcal{D}}$,

$$(9) \quad |x(t, x_0, d)|_{\mathcal{A}} < \varepsilon \quad \text{whenever } |x_0|_{\mathcal{A}} < r \text{ and } t \geq T_r(\varepsilon).$$

Proof. Sufficiency is clear. Now we show the necessity part. For any $r, \varepsilon > 0$, let

$$(10) \quad A_{r, \varepsilon} \stackrel{\text{def}}{=} \{T \geq 0 : \forall |x_0|_{\mathcal{A}} < r, \forall t \geq T, \forall d \in \mathcal{M}_{\mathcal{D}}, |x(t, x_0, d)|_{\mathcal{A}} < \varepsilon\} \subseteq \mathbb{R}_{\geq 0}.$$

Then from the assumptions, $A_{r, \varepsilon} \neq \emptyset$ for any $r, \varepsilon > 0$. Moreover,

$$A_{r, \varepsilon_1} \subseteq A_{r, \varepsilon_2}, \text{ if } \varepsilon_1 \leq \varepsilon_2, \text{ and } A_{r_2, \varepsilon} \subseteq A_{r_1, \varepsilon}, \text{ if } r_1 \leq r_2.$$

Now define $\bar{T}_r(\varepsilon) \stackrel{\text{def}}{=} \inf A_{r, \varepsilon}$. Then $\bar{T}_r(\varepsilon) < \infty$, for any $r, \varepsilon > 0$, and it satisfies

$$\bar{T}_r(\varepsilon_1) \geq \bar{T}_r(\varepsilon_2), \text{ if } \varepsilon_1 \leq \varepsilon_2, \text{ and } \bar{T}_{r_1}(\varepsilon) \leq \bar{T}_{r_2}(\varepsilon), \text{ if } r_1 \leq r_2.$$

So we can define for any $r, \varepsilon > 0$,

$$(11) \quad \tilde{T}_r(\varepsilon) \stackrel{\text{def}}{=} \frac{2}{\varepsilon} \int_{\varepsilon/2}^{\varepsilon} \bar{T}_r(s) ds.$$

Since $\bar{T}_r(\cdot)$ is decreasing, $\tilde{T}_r(\cdot)$ is well defined and is locally absolutely continuous. Also

$$(12) \quad \tilde{T}_r(\varepsilon) \geq \frac{2}{\varepsilon} \bar{T}_r(\varepsilon) \int_{\varepsilon/2}^{\varepsilon} ds = \bar{T}_r(\varepsilon).$$

Furthermore,

$$(13) \quad \begin{aligned} \frac{d\tilde{T}_r(\varepsilon)}{d\varepsilon} &= -\frac{2}{\varepsilon^2} \int_{\varepsilon/2}^{\varepsilon} \bar{T}_r(s) ds + \frac{2}{\varepsilon} \left(\bar{T}_r(\varepsilon) - \frac{1}{2} \bar{T}_r\left(\frac{\varepsilon}{2}\right) \right) \\ &= \frac{1}{\varepsilon} \left[\bar{T}_r(\varepsilon) - \frac{2}{\varepsilon} \int_{\varepsilon/2}^{\varepsilon} \bar{T}_r(s) ds \right] + \frac{1}{\varepsilon} \left[\bar{T}_r(\varepsilon) - \bar{T}_r\left(\frac{\varepsilon}{2}\right) \right] \\ &= \frac{1}{\varepsilon} \left[\bar{T}_r(\varepsilon) - \tilde{T}_r(\varepsilon) \right] + \frac{1}{\varepsilon} \left[\bar{T}_r(\varepsilon) - \bar{T}_r\left(\frac{\varepsilon}{2}\right) \right] \leq 0, \text{ a.e.,} \end{aligned}$$

hence $\tilde{T}_r(\cdot)$ decreases (not necessarily strictly). Since $\bar{T}_{(\cdot)}(\varepsilon)$ increases, from the definition, $\tilde{T}_{(\cdot)}(\varepsilon)$ also increases. Finally, define

$$(14) \quad T_r(\varepsilon) \stackrel{\text{def}}{=} \tilde{T}_r(\varepsilon) + \frac{r}{\varepsilon}$$

Then it follows that

- for any fixed r , $T_r(\cdot)$ is continuous, maps $\mathbb{R}_{>0} \xrightarrow{\text{onto}} \mathbb{R}_{>0}$, and is strictly decreasing;
- for any fixed ε , $T_r(\varepsilon)$ is increasing as r increases, and $\lim_{r \rightarrow \infty} T_r(\varepsilon) = \infty$.

So the only thing left to be shown is that T_r defined by (14) satisfies (9). To do this, pick any x_0 and t with $|x_0|_{\mathcal{A}} < r$ and $t \geq T_r(\varepsilon)$. Then

$$t \geq T_r(\varepsilon) > \tilde{T}_r(\varepsilon) \geq \bar{T}_r(\varepsilon).$$

Hence, by the definition of $\bar{T}_r(\varepsilon)$, $|x(t, x_0, d)|_{\mathcal{A}} < \varepsilon$, as claimed. \square

3.1. Proof of Characterization via Decay Estimate. We now provide a proof of Proposition 2.5.

[\Leftarrow] Assume that there exists a \mathcal{KL} -function β such that (6) holds. Let

$$c_1 \stackrel{\text{def}}{=} \sup \beta(\cdot, 0) \leq \infty,$$

and choose $\delta(\cdot)$ to be any \mathcal{K}_{∞} -function with

$$\delta(\varepsilon) \leq \bar{\beta}^{-1}(\varepsilon), \text{ any } 0 \leq \varepsilon < c_1,$$

where $\bar{\beta}^{-1}$ denotes the inverse function of $\bar{\beta}(\cdot) \stackrel{\text{def}}{=} \beta(\cdot, 0)$. (If $c_1 = \infty$, we can simply choose $\delta(\varepsilon) \stackrel{\text{def}}{=} \bar{\beta}^{-1}(\varepsilon)$.) Clearly $\delta(\varepsilon)$ is the desired \mathcal{K}_∞ -function for the uniform stability property.

The uniform attraction property follows from the fact that for every fixed r , $\lim_{t \rightarrow \infty} \beta(r, t) = 0$.

[\implies] Assume that (1) is UGAS with respect to the closed set \mathcal{A} , and let δ be as in the definition. Let $\varphi(\cdot)$ be the \mathcal{K} -function $\delta^{-1}(\cdot)$. As mentioned in Remark 2.4, it follows that $|x(t, x_0, d)|_{\mathcal{A}} \leq \varphi(|x_0|_{\mathcal{A}})$ for any $x_0 \in \mathbb{R}^n$, any $t \geq 0$, and any $d \in \mathcal{M}_{\mathcal{D}}$.

Let $\{T_r\}_{r \in (0, \infty)}$ be as in Lemma 3.1, and for each $r \in (0, \infty)$ denote $\psi_r \stackrel{\text{def}}{=} T_r^{-1}$. Then, for each $r \in (0, \infty)$, $\psi_r : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ is again continuous, onto, and strictly decreasing. We also write $\psi_r(0) = +\infty$, which is consistent with that fact that

$$\lim_{t \rightarrow 0^+} \psi_r(t) = +\infty.$$

(Note: The property that $T_{(\cdot)}(t)$ increases to ∞ is not needed here.)

Claim: For any $|x_0|_{\mathcal{A}} < r$, any $t \geq 0$ and any $d \in \mathcal{M}_{\mathcal{D}}$, $|x(t, x_0, d)|_{\mathcal{A}} \leq \psi_r(t)$.

Proof: It follows from the definition of the maps T_r that, for any $r, \varepsilon > 0$, and for any $d \in \mathcal{M}_{\mathcal{D}}$,

$$|x_0|_{\mathcal{A}} < r, \quad t \geq T_r(\varepsilon) \implies |x(t, x_0, d)|_{\mathcal{A}} < \varepsilon.$$

As $t = T_r(\psi_r(t))$ if $t > 0$, we have, for any such x_0 and d ,

$$(15) \quad |x(t, x_0, d)|_{\mathcal{A}} < \psi_r(t), \quad \forall t > 0.$$

The claim follows by combining (15) and the fact that $\psi_r(0) = +\infty$.

Now for any $s \geq 0$ and $t \geq 0$, let

$$(16) \quad \bar{\psi}(s, t) \stackrel{\text{def}}{=} \min \left\{ \varphi(s), \inf_{r \in (s, \infty)} \psi_r(t) \right\}.$$

Because of the definition of φ and the above claim, we have, for each $x_0, d \in \mathcal{M}_{\mathcal{D}}$, and $t \geq 0$:

$$(17) \quad |x(t, x_0, d)|_{\mathcal{A}} \leq \bar{\psi}(|x_0|_{\mathcal{A}}, t).$$

If $\bar{\psi}$ would be of class \mathcal{KL} , we would be done. This may not be the case, so we next majorize $\bar{\psi}$ by such a function.

By its definition, for any fixed t , $\bar{\psi}(\cdot, t)$ is an increasing function (not necessarily strictly). Also because for any fixed $r \in (0, \infty)$, $\psi_r(t)$ decreases to 0 (this follows from the fact that $\psi_r : \mathbb{R}_{>0} \xrightarrow{\text{onto}} \mathbb{R}_{>0}$ is continuous and strictly decreasing), it follows that

for any fixed s , $\bar{\psi}(s, t)$ decreases to 0 as $t \rightarrow \infty$.

Next we construct a function $\tilde{\psi} : \mathbb{R}_{[0, \infty)} \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ with the following properties:

- for any fixed $t \geq 0$, $\tilde{\psi}(\cdot, t)$ is continuous and strictly increasing;
- for any fixed $s \geq 0$, $\tilde{\psi}(s, t)$ decreases to 0 as $t \rightarrow \infty$;
- $\tilde{\psi}(s, t) \geq \bar{\psi}(s, t)$.

Such a function $\tilde{\psi}$ always exists; for instance, it can be obtained as follows. Define first

$$(18) \quad \hat{\psi}(s, t) \stackrel{\text{def}}{=} \int_s^{s+1} \bar{\psi}(\varepsilon, t) d\varepsilon.$$

Then $\hat{\psi}(\cdot, t)$ is an absolutely continuous function on every compact subset of $\mathbb{R}_{\geq 0}$, and it satisfies

$$\hat{\psi}(s, t) \geq \bar{\psi}(s, t) \int_s^{s+1} d\varepsilon = \bar{\psi}(s, t).$$

It follows that

$$\frac{\partial \hat{\psi}(s, t)}{\partial s} = \bar{\psi}(s+1, t) - \bar{\psi}(s, t) \geq 0, \text{ a.e.},$$

and hence $\hat{\psi}(\cdot, t)$ is increasing. Also since for any fixed s , $\bar{\psi}(s, \cdot)$ decreases, so does $\hat{\psi}(s, \cdot)$. Note that

$$\bar{\psi}(s, t) \leq \bar{\psi}(s, 0) = \min \left\{ \inf_{r \in (s, \infty)} \psi_r(0), \varphi(s) \right\} = \varphi(s),$$

(recall that $\psi_r(0) = +\infty$), so by the Lebesgue dominated convergence theorem, for any fixed $s \geq 0$,

$$\lim_{t \rightarrow \infty} \hat{\psi}(s, t) = \int_s^{s+1} \lim_{t \rightarrow \infty} \bar{\psi}(\varepsilon, t) d\varepsilon = 0.$$

Now we see that the function $\hat{\psi}(s, t)$ satisfies all of the requirements for $\tilde{\psi}(s, t)$ except possibly for the strictly increasing property. We define $\tilde{\psi}$ as follows:

$$\tilde{\psi}(s, t) \stackrel{\text{def}}{=} \hat{\psi}(s, t) + \frac{s}{(s+1)(t+1)}.$$

Clearly it satisfies all the desired properties.

Finally, define

$$\beta(s, t) \stackrel{\text{def}}{=} \sqrt{\varphi(s)} \sqrt{\tilde{\psi}(s, t)}.$$

Then it follows that $\beta(s, t)$ is a \mathcal{KL} -function, and, for all x_0, t, d :

$$|x(t, x_0, d)|_{\mathcal{A}} \leq \sqrt{\varphi(|x_0|_{\mathcal{A}})} \sqrt{\tilde{\psi}(|x_0|_{\mathcal{A}}, t)} \leq \beta(|x_0|_{\mathcal{A}}, t),$$

which concludes the proof of the Proposition.

4. Some Preliminaries about Lyapunov Functions. In this section we provide some technical results about set Lyapunov functions. A lemma on differential inequalities is also given, for later reference.

REMARK 4.1. One may assume in Definition 2.6 that all of $\alpha_1, \alpha_2, \alpha_3$ are smooth in $(0, +\infty)$ and of class \mathcal{K}_∞ . For α_1 and α_2 , this is proved simply by finding two functions $\tilde{\alpha}_1, \tilde{\alpha}_2$ in \mathcal{K}_∞ , smooth in $(0, +\infty)$ so that

$$\tilde{\alpha}_1(s) \leq \alpha_1(s) \leq \alpha_2(s) \leq \tilde{\alpha}_2(s), \text{ for all } s.$$

For α_3 , a new Lyapunov function W and a function $\tilde{\alpha}_3$ which satisfies (8) with respect to W , but is smooth in $(0, +\infty)$ and of class \mathcal{K}_∞ , can be constructed as follows. First, pick $\tilde{\alpha}_3$ to be any \mathcal{K}_∞ -function, smooth in $(0, +\infty)$, such that

$$\tilde{\alpha}_3(s) \leq s\alpha_3(s), \forall s \in [0, \alpha_1^{-1}(1)].$$

This is possible since α_3 is positive definite. Then let

$$\gamma : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}_{\geq 0}$$

be a \mathcal{K}_∞ -function, smooth in $(0, +\infty)$, such that

- $\gamma(r) \geq \alpha_1^{-1}(r)$ for all $r \in [0, 1]$;
- $\gamma(r) > \frac{\tilde{\alpha}_3(\alpha_1^{-1}(r))}{\alpha_3(\alpha_1^{-1}(r))}$ for all $r > 1$.

Now define $\beta(s) \stackrel{\text{def}}{=} \int_0^s \gamma(r) dr$. Note that β is a \mathcal{K}_∞ -function, smooth in $(0, +\infty)$. Let $W(\xi) \stackrel{\text{def}}{=} \beta(V(\xi))$. This is smooth on $\mathbb{R}^n \setminus \mathcal{A}$, and $\beta \circ \alpha_1, \beta \circ \alpha_2$ bound W as in equation (7). Moreover,

$$\beta'(V(\xi)) = \gamma(V(\xi)) \geq \gamma(\alpha_1(|\xi|_{\mathcal{A}})),$$

so

$$(19) \quad L_{f_d} W(\xi) = \beta'(V(\xi)) L_{f_d} V(\xi) \leq -\gamma(\alpha_1(|\xi|_{\mathcal{A}})) \alpha_3(|\xi|_{\mathcal{A}}).$$

We claim that this is bounded by $-\tilde{\alpha}_3(|\xi|_{\mathcal{A}})$. Indeed, if $s \stackrel{\text{def}}{=} |\xi|_{\mathcal{A}} \leq \alpha_1^{-1}(1)$, then from the first item above and the definition of $\tilde{\alpha}_3$,

$$\gamma(\alpha_1(s)) \geq s \geq \frac{\tilde{\alpha}_3(s)}{\alpha_3(s)};$$

if instead $s > \alpha_1^{-1}(1)$, then from the second item, also

$$\gamma(\alpha_1(s)) \geq \frac{\tilde{\alpha}_3(s)}{\alpha_3(s)}.$$

In either case, $\gamma(\alpha_1(s)) \alpha_3(s) \geq \tilde{\alpha}_3(s)$, as desired. From now on, whenever necessary, we assume that $\alpha_1, \alpha_2, \alpha_3$ are \mathcal{K}_∞ -functions, smooth in $(0, +\infty)$. \square

4.1. Smoothing of Lyapunov Functions. When dealing with control system design, one often needs to know that V can be taken to be globally smooth, rather than just smooth outside of \mathcal{A} .

PROPOSITION 4.2. *If there is a Lyapunov function for (1) with respect to \mathcal{A} , then there is also a smooth such Lyapunov function.*

The proof relies on constructing a smooth function of the form $W = \beta \circ V$, where

$$\beta : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}_{\geq 0}$$

is built using a partition of unity.

Again let $\mathcal{A} \subseteq \mathbb{R}^n$ be nonempty and closed. For a multi-index $\varrho = (\varrho_1, \varrho_2, \dots, \varrho_n)$, we use $|\varrho|$ to denote $\sum_{i=1}^n \varrho_i$. The following regularization result will be needed; it generalizes to arbitrary \mathcal{A} the analogous (but simpler, due to compactness) result for equilibria given in [13, Theorem 6].

LEMMA 4.3. *Assume that $V : \mathbb{R}^n \longrightarrow \mathbb{R}_{\geq 0}$ is C^0 , the restriction $V|_{\mathbb{R}^n \setminus \mathcal{A}}$ is C^∞ , and also $V|_{\mathcal{A}} = 0$, $V|_{\mathbb{R}^n \setminus \mathcal{A}} > 0$. Then there exists a \mathcal{K}_∞ -function β , smooth on $(0, \infty)$ and so that $\beta^{(i)}(t) \rightarrow 0$ as $t \rightarrow 0^+$ for each $i = 0, 1, \dots$ and having $\beta'(t) > 0, \forall t > 0$, such that*

$$W \stackrel{\text{def}}{=} \beta \circ V$$

is a C^∞ function on all of \mathbb{R}^n .

Proof. Let K_1, K_2, \dots , be compact subsets of \mathbb{R}^n such that $\mathcal{A} \subseteq \bigcup_{i=1}^\infty \text{int}(K_i)$. For any $k \geq 1$, let

$$I_k \stackrel{\text{def}}{=} \left(\frac{1}{k+2}, \frac{1}{k} \right) \subseteq \mathbb{R}$$

and $I_0 \stackrel{\text{def}}{=} I_1$. Pick for any $k \geq 1$, a smooth (C^∞) function $\gamma_k : \mathbb{R}_{>0} \rightarrow [0, 1]$ satisfying

- $\gamma_k(t) = 0$ if $t \notin I_k$; and
- $\gamma_k(t) > 0$ if $t \in I_k$.

Define for any $k \geq 1$,

$$\mathcal{G}_k \stackrel{\text{def}}{=} \left\{ x \in \mathbb{R}^n : x \in \bigcup_{i=1}^k K_i, V(x) \in \text{clos } I_k \right\}.$$

Then \mathcal{G}_k is compact (because of compactness of the sets K_i and continuity of V). Observe that each derivative $\gamma_k^{(i)}$ has a compact support included in $\text{clos } I_k$, so it is bounded. For each $k = 1, 2, \dots$, let $c_k \in \mathbb{R}$ satisfy

1. $c_k \geq 1$;
2. $c_k \geq |(D^\varrho V)(x)|$ for any multi-index $|\varrho| \leq k$ and any $x \in \mathcal{G}_k$; and
3. $c_k \geq |\gamma_k^{(i)}(t)|$, for any $i \leq k$ and any $t \in \mathbb{R}_{>0}$.

Choose the sequence d_k to satisfy

$$(20) \quad 0 < d_k < \frac{1}{2^k (k+1)! c_k^k}, \quad k = 1, 2, \dots$$

Let $\alpha : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be a C^∞ function such that $\alpha \equiv 0$ on $\left[0, \frac{1}{3}\right]$ and $\alpha \geq 1$ on $\left[\frac{1}{2}, \infty\right)$. Define $\gamma(0) \stackrel{\text{def}}{=} 0$ and

$$(21) \quad \gamma(t) \stackrel{\text{def}}{=} \sum_{k=1}^{\infty} d_k \gamma_k(t) + \alpha(t), \quad \forall t > 0.$$

Notice that for any $t \in (0, 1)$, if $k \stackrel{\text{def}}{=} \left\lfloor \frac{1}{t} \right\rfloor \geq 1$ denotes the largest integer $\leq \frac{1}{t}$, then $t \in I_{k-1}$, and

$$t \notin I_j \quad \text{if } j \neq k, k-1.$$

Hence the sum in (21) at most consists of three terms (for $t \geq 1$ the sum is just $\gamma = \alpha$), and so γ is C^∞ at each $t \in (0, \infty)$.

Claim: For any $i \geq 0$, $\lim_{t \rightarrow 0^+} \gamma^{(i)}(t) = 0$.

Proof: Fix any $i \geq 0$. Given any $\varepsilon > 0$, let $k_0 \in \mathbb{Z}$ be such that $\varepsilon > \frac{1}{k_0} > 0$. Let

$$T \stackrel{\text{def}}{=} \min \left\{ \frac{1}{k_0}, \frac{1}{i+1}, \frac{1}{3} \right\}.$$

We will show that $t \in (0, T) \implies |\gamma^{(i)}(t)| < \varepsilon$. Indeed, as $0 < t < \min \left\{ \frac{1}{k_0}, \frac{1}{i+1}, \frac{1}{3} \right\}$, it follows that $k \stackrel{\text{def}}{=} \left\lfloor \frac{1}{t} \right\rfloor \geq \max\{i+1, k_0, 3\}$. So

$$\gamma^{(i)}(t) \leq d_{k-1} \gamma_{k-1}^{(i)}(t) + d_k \gamma_k^{(i)}(t),$$

and noticing that

$$i \leq k-1 < k \implies c_k \geq \left| \gamma_k^{(i)}(t) \right|, \quad c_{k-1} \geq \left| \gamma_{k-1}^{(i)}(t) \right|,$$

we have

$$\left| \gamma^{(i)}(t) \right| \leq d_{k-1} c_{k-1} + d_k c_k \leq \frac{1}{2k!} + \frac{1}{2(k+1)!} < \frac{1}{k!} < \frac{1}{k} \leq \frac{1}{k_0} < \varepsilon,$$

as wanted.

Note also that if $t \geq \frac{1}{2}$, then $\gamma(t) \geq \alpha(t) \geq 1 > 0$; and if $t \in \left(0, \frac{1}{2}\right)$, then $\gamma(t) \geq d_{k-1} \gamma_{k-1}(t) > 0$ with $k \stackrel{\text{def}}{=} \left\lfloor \frac{1}{t} \right\rfloor \geq 2$, so the function

$$(22) \quad \beta(t) \stackrel{\text{def}}{=} \int_0^t \gamma(s) ds$$

is also a \mathcal{K}_∞ -function, smooth on $(0, \infty)$. Furthermore, β satisfies $\beta^{(i)}(t) \rightarrow 0$ as $t \rightarrow 0^+$ for each $i = 0, 1, \dots$

Finally, we show that $W = \beta \circ V$ is C^∞ . For this, it is enough to show that $D^{e_0} W(x_n) \rightarrow 0$ as $x_n \rightarrow \bar{x} \in \partial \mathcal{A}$, for each multi-index e_0 and each sequence $\{x_n\} \subseteq \mathbb{R}^n \setminus \mathcal{A}$ converging to a point \bar{x} in the boundary of \mathcal{A} . (In general, see e.g. [4] (p. 52), if $\mathcal{A} \subseteq \mathbb{R}^n$ is closed and $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies that $\varphi|_{\mathcal{A}} = 0$, $\varphi|_{\mathbb{R}^n \setminus \mathcal{A}}$ is C^∞ , and for

each boundary point a of \mathcal{A} and all multi-indices $\varrho = (\varrho_1, \varrho_2, \dots, \varrho_n)$, it holds that $\lim_{\substack{x \rightarrow a \\ x \notin \mathcal{A}}} D^\varrho \varphi(x) = 0$, then φ is C^∞ on \mathbb{R}^n .)

Pick one such ϱ_0 and any sequence $\{x_n\}$ with $x_n \rightarrow \bar{x} \in \partial\mathcal{A}$. If $|\varrho_0| = 0$, one only needs to show that $W(x_n) \rightarrow 0$, which follows easily from the facts that $\beta \in \mathcal{K}_\infty$ and $V(x_n) \rightarrow 0$. So from now on, we can assume that $|\varrho_0| \stackrel{\text{def}}{=} i \geq 1$. As $\mathcal{A} \subseteq \cup_{j=0}^\infty \text{int } K_j$, $\bar{x} \in \text{int } K_l$ for some l , and without loss of generality we may assume that there is some fixed l so that

$$x_n \in K_l, \quad \text{for all } n.$$

Pick any $\varepsilon > 0$. We will show that there exists some N such that

$$n > N \implies |D^{\varrho_0} W(x_n)| < \varepsilon.$$

Let $k \in \mathbb{Z}$ be so that

$$k > \max \left\{ i, \log_2 \left(\frac{1}{\varepsilon} \right), l \right\}$$

and let $T \in \left(0, \frac{1}{3}\right)$ be such that $T < \frac{1}{k+2}$. Observe that if $t < T$, then $t \notin I_1 \cup \dots \cup I_k$.

As V is C^0 everywhere, $V = 0$ at \mathcal{A} , $V(x_n) \rightarrow V(\bar{x}) = 0$. So there exists N such that $V(x_n) < T$ whenever $n > N$. Fix an N like this. Then for any $n > N$,

$$\gamma_s^{(j)}(V(x_n)) = 0, \quad \forall j, \forall s = 1, 2, \dots, k,$$

(since γ_s vanishes outside I_s). Pick any $j \in \mathbb{N}$ with $j \leq i$, any $h \in \mathbb{N}$ with $h \leq i$, and $\varrho_1, \dots, \varrho_h$ multi-indices such that $|\varrho_\mu| \leq i$, $\forall \mu = 1, \dots, h$. Then for any $q \in \mathbb{N}$ with $q > k$, by the way we chose c_k ,

$$|\gamma_q^{(j)}(V(x_n))| \leq c_q,$$

since $q > k > i \geq j$. Also, if $V(x_n) \in I_q$, then again by the properties of the sequence c_k ,

$$|D^{\varrho_\mu} V(x_n)| \leq c_q,$$

(since $q > k > l$ and $x_n \in K_l$ imply $x_n \in K_1 \cup \dots \cup K_q$, and $|\varrho_\mu| \leq i < k < q$). Therefore, for such q , if $V(x_n) \in I_q$,

$$(23) \quad |\gamma_q^{(j)}(V(x_n))| |D^{\varrho_1} V(x_n)| \cdots |D^{\varrho_h} V(x_n)| \leq c_q^{h+1} \leq c_q^{i+1} < c_q^i.$$

If instead it would be the case that $V(x_n) \notin I_q$, then $\gamma_q^{(j)}(V(x_n)) = 0$, and hence the inequality (23) still holds. Since

$$\gamma^{(j)}(V(x_n)) = \sum_{q=k+1}^{\infty} d_q \gamma_q^{(j)}(V(x_n)),$$

we also have

$$(24) \quad \begin{aligned} & |\gamma^{(j)}(V(x_n))| |D^{\varrho_1} V(x_n)| \cdots |D^{\varrho_h} V(x_n)| \leq \sum_{q=k+1}^{\infty} d_q c_q^i < \sum_{q=k+1}^{\infty} \frac{1}{2^q (q+1)!} \\ & < \left(\sum_{q=k+1}^{\infty} \frac{1}{2^q} \right) \frac{1}{(k+1)!} = \frac{1}{2^k (k+1)!} < \frac{\varepsilon}{(k+1)!}. \end{aligned}$$

Now observe that

$$(D^{e_0} W)(x) = (D^{e_0}(\beta \circ V))(x)$$

is a sum of $\leq i!$ terms (recall $0 < i = |\varrho_0|$), each of which is of the form

$$\beta^{(p)}(V(x)) (D^{e_1} V)(x) \cdots (D^{e_h} V)(x),$$

where $0 < p \leq i$, $h \leq i$, and each $|\varrho_\mu| \leq i$. Each

$$\beta^{(p)}(V(x)) = \gamma^{(j)}(V(x)), \quad j = p - 1 \leq i - 1,$$

so (24) applies, and we conclude

$$|(D^{e_0} W)(x_n)| \leq i! \frac{\varepsilon}{(k+1)!} < \varepsilon,$$

(since $k > i$.) \square

Now let us return to the proof of the Proposition 4.2.

Proof of Proposition 4.2. Assume \mathcal{A} , V and $\alpha_1, \alpha_2, \alpha_3$ are as defined in Definition 2.6. Let β, W be as in Lemma 4.3. We show that W is a smooth Lyapunov function as required.

Let $\hat{\alpha}_i \stackrel{\text{def}}{=} \beta \circ \alpha_i$, $i = 1, 2$. These are again \mathcal{K}_∞ -functions, and they satisfy

$$\hat{\alpha}_1(|\xi|_{\mathcal{A}}) \leq W(\xi) \leq \hat{\alpha}_2(|\xi|_{\mathcal{A}}).$$

We define, for $s > 0$:

$$\check{\beta}(s) \stackrel{\text{def}}{=} \min_{t \in [\alpha_1(s), \alpha_2(s)]} \beta'(t) > 0.$$

Let also $\check{\beta}(0) \stackrel{\text{def}}{=} 0$. Define $\hat{\alpha}_3(s) \stackrel{\text{def}}{=} \check{\beta}(s)\alpha_3(s)$. Then $\hat{\alpha}_3$ is a continuous positive definite function. Also, for any $\xi \in \mathbb{R}^n \setminus \mathcal{A}$,

$$\begin{aligned} L_{f_d} W(\xi) &= \beta'(V(\xi)) L_{f_d} V(\xi) \leq -\beta'(V(\xi)) \alpha_3(|\xi|_{\mathcal{A}}) \\ &\leq -\check{\beta}(|\xi|_{\mathcal{A}}) \alpha_3(|\xi|_{\mathcal{A}}) = -\hat{\alpha}_3(|\xi|_{\mathcal{A}}), \end{aligned}$$

which concludes the proof of the Proposition. \blacksquare

4.2. A Useful Estimate. The following lemma establishes a useful comparison principle.

LEMMA 4.4. *For each continuous and positive definite function α , there exists a \mathcal{KL} -function $\beta_\alpha(s, t)$ with the following property: if $y(\cdot)$ is any (locally) absolutely continuous function defined for $t \geq 0$ and with $y(t) \geq 0$ for all t , and $y(\cdot)$ satisfies the differential inequality*

$$(25) \quad \dot{y}(t) \leq -\alpha(y(t)), \quad \text{for almost all } t$$

with $y(0) = y_0 \geq 0$, then it holds that

$$y(t) \leq \beta_\alpha(y_0, t)$$

for all $t \geq 0$.

Proof. Define for any $s > 0$, $\eta(s) \stackrel{\text{def}}{=} -\int_1^s \frac{dr}{\alpha(r)}$. This is a strictly decreasing differentiable function on $(0, \infty)$. Without loss of generality, we will assume that $\lim_{s \rightarrow 0^+} \eta(s) = +\infty$. If this were not the case, we could consider instead the following function:

$$\bar{\alpha}(s) \stackrel{\text{def}}{=} \begin{cases} \min\{s, \alpha(s)\}, & \text{if } 0 \leq s < 1, \\ \alpha(s), & \text{if } s \geq 1. \end{cases}$$

This function is again continuous, positive definite, satisfies $\bar{\alpha}(s) \leq \alpha(s)$ for any $s \geq 0$, and

$$\lim_{s \rightarrow 0^+} \int_s^1 \frac{dr}{\bar{\alpha}(r)} \geq \int_s^1 \frac{dr}{r} = +\infty.$$

Moreover, if $\dot{y}(t) \leq -\alpha(y(t))$ then also $\dot{y}(t) \leq -\bar{\alpha}(y(t))$, so $\beta_{\bar{\alpha}}$ could be used to bound solutions.

Let

$$0 < a \stackrel{\text{def}}{=} - \lim_{s \rightarrow +\infty} \eta(s).$$

Then the range of η , and hence also the domain of η^{-1} , is the open interval $(-a, \infty)$. (We allow the possibility that $a = \infty$.) For $(s, t) \in \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}$, define

$$\beta_{\alpha}(s, t) \stackrel{\text{def}}{=} \begin{cases} 0, & \text{if } s = 0, \\ \eta^{-1}(\eta(s) + t), & \text{if } s > 0. \end{cases}$$

We claim that for any $y(\cdot)$ satisfying the conditions in the Lemma,

$$(26) \quad y(t) \leq \beta_{\alpha}(y_0, t), \quad \text{for all } t \geq 0.$$

As $\dot{y}(t) \leq -\alpha(y(t))$, it follows that $y(t)$ is nonincreasing, and if $y(t_0) = 0$ for some $t_0 \geq 0$, then $y(t) \equiv 0$, $\forall t \geq t_0$. Without loss of generality, assume that $y_0 > 0$. Let

$$t_0 \stackrel{\text{def}}{=} \inf\{t : y(t) = 0\} \leq +\infty.$$

It is enough to show (26) holds for $t \in [0, t_0)$.

As η is strictly decreasing, we only need to show that $\eta(y(t)) \geq \eta(y_0) + t$, that is,

$$- \int_1^{y(t)} \frac{dr}{\alpha(r)} \geq - \int_1^{y_0} \frac{dr}{\alpha(r)} + t,$$

which is equivalent to

$$(27) \quad \int_{y(t)}^{y_0} \frac{dr}{\alpha(r)} \geq t.$$

From (25), one sees that

$$\int_0^t \frac{\dot{y}(\tau)}{\alpha(y(\tau))} d\tau \leq - \int_0^t d\tau = -t.$$

Changing variables in the integral, this gives (27).

It only remains to show that β_{α} is of class \mathcal{KL} . The function β_{α} is continuous since both η and η^{-1} are continuous in their domains, and $\lim_{r \rightarrow \infty} \eta^{-1}(r) = 0$. It is strictly increasing in s for each fixed t since since both η and η^{-1} are strictly decreasing. Finally, $\beta_{\alpha}(s, t) \rightarrow 0$ as $t \rightarrow \infty$ by construction. So β_{α} is a \mathcal{KL} -function. \square

5. Some Properties of Complete Systems. We need to first establish some technical properties that hold for complete systems, and in particular a Lipschitz continuity fact.

For each $\xi \in \mathbb{R}^n$ and $T > 0$, let

$$\mathcal{R}^T(\xi) \stackrel{\text{def}}{=} \{\eta : \eta = x(T, \xi, d), d \in \mathcal{M}_{\mathcal{D}}\}.$$

This is the reachable set of (1) from ξ at time T . We use $\mathcal{R}^{\leq T}(\xi)$ to denote $\bigcup_{0 \leq t \leq T} \mathcal{R}^t(\xi)$. If S is a subset of \mathbb{R}^n ,

we write

$$\mathcal{R}^T(S) \stackrel{\text{def}}{=} \bigcup_{\xi \in S} \mathcal{R}^T(\xi), \quad \mathcal{R}^{\leq T}(S) \stackrel{\text{def}}{=} \bigcup_{\xi \in S} \mathcal{R}^{\leq T}(\xi).$$

In what follows we use \bar{S} to denote the closure of S for any subset S of \mathbb{R}^n .

PROPOSITION 5.1. *Assume that (1) is forward complete. Then for any compact subset K of \mathbb{R}^n and any $T > 0$, the set $\overline{\mathcal{R}^{\leq T}(K)}$ is compact.*

To prove Proposition 5.1, we first need to make a couple of technical observations.

LEMMA 5.2. *Let K be a compact subset of \mathbb{R}^n and let $T > 0$. Then the set $\overline{\mathcal{R}^{\leq T}(K)}$ is compact if and only if $\overline{\mathcal{R}^{\leq T}(\xi)}$ is compact for each $\xi \in K$.*

Proof. It is clear that the compactness of $\overline{\mathcal{R}^{\leq T}(K)}$ implies the compactness of $\overline{\mathcal{R}^{\leq T}(\xi)}$ for any $\xi \in K$.

Now assume, for $T > 0$ and a compact set K , that $\overline{\mathcal{R}^{\leq T}(\xi)}$ is compact for each $\xi \in K$. Pick any $\xi \in K$, and let $\mathcal{U} = \{\eta : d(\eta, \overline{\mathcal{R}^{\leq T}(\xi)}) < 1\}$. Then $\bar{\mathcal{U}}$ is compact. Let C be a Lipschitz constant for f with respect to x on $\bar{\mathcal{U}}$, and let $r = e^{-CT}$. For each $d \in \mathcal{M}_{\mathcal{D}}$ and each η with $|\eta - \xi| < r$, let $\tilde{t} = \inf\{t \geq 0 : |x(t, \eta, d) - x(t, \xi, d)| \geq 1\}$. Then, using Gronwall's Lemma, one can show that $\tilde{t} \geq T$, from which it follows that

$$\mathcal{R}^{\leq T}(\eta) \subseteq \bar{\mathcal{U}}, \quad \forall |\eta - \xi| < r.$$

Thus, for each $\xi \in K$, there is a neighborhood \mathcal{V}_{ξ} of ξ such that $\overline{\mathcal{R}^{\leq T}(\mathcal{V}_{\xi})}$ is compact. By compactness of K , it follows that $\overline{\mathcal{R}^{\leq T}(K)}$ is compact. \square

LEMMA 5.3. *For any subset S of \mathbb{R}^n and any $T > 0$,*

$$\mathcal{R}^T(\bar{S}) \subseteq \overline{\mathcal{R}^T(S)}, \quad \mathcal{R}^{\leq T}(\bar{S}) \subseteq \overline{\mathcal{R}^{\leq T}(S)}.$$

In particular, $\overline{\mathcal{R}^{\leq T}(\bar{S})} = \overline{\mathcal{R}^T(S)}$.

Proof. The first conclusion follows from the continuity of solutions on initial states; see [26], Theorem 1. The second is immediate from there. \square

We now return to the proof of Proposition 5.1. By Lemma 5.2, it is enough to show that $\overline{\mathcal{R}^{\leq T}(\xi)}$ is compact for each $\xi \in \mathbb{R}^n$ and each $T > 0$. Pick any $\xi_0 \in \mathbb{R}^n$, and let

$$\tau = \sup\{T \geq 0 : \overline{\mathcal{R}^{\leq T}(\xi_0)} \text{ is compact}\}.$$

Note that $\tau > 0$. This is because $|x(t, \xi_0, d) - \xi_0| \leq 1$ for any $0 \leq t < 1/M$ and any $d \in \mathcal{M}_{\mathcal{D}}$, where

$$M = \max\{|f(\xi, \mathbf{d})| : |\xi - \xi_0| \leq 1, \mathbf{d} \in \mathcal{D}\}.$$

We must show that $\tau = \infty$.

Assume that $\tau < \infty$. Using the same argument used above, one can show that if $\overline{\mathcal{R}^{\leq t}(\xi_0)}$ is compact for some $t > 0$ then there is some $\delta > 0$ such that $\overline{\mathcal{R}^{\leq (t+\delta)}(\xi_0)}$ is compact. From here it follows that $\overline{\mathcal{R}^{\leq \tau}(\xi_0)}$ is not compact. By definition, $\overline{\mathcal{R}^{\leq t}(\xi_0)}$ is compact for any $t < \tau$.

Let $\tau_1 = \tau/2$. Then there is some $\eta_1 \in \overline{\mathcal{R}^{\tau_1}(\xi_0)}$ such that $\overline{\mathcal{R}^{\leq (\tau-\tau_1)}(\eta_1)}$ is not compact; otherwise, by Lemma 5.2, $\overline{\mathcal{R}^{\leq (\tau-\tau_1)}(\overline{\mathcal{R}^{\tau_1}(\xi_0)})}$ would be compact. This, in turn, would imply that $\overline{\mathcal{R}^{\leq \tau}(\xi_0)}$ is compact, since

$$\mathcal{R}^{\leq \tau}(\xi_0) \subseteq \mathcal{R}^{\leq \tau_1}(\xi_0) \cup \mathcal{R}^{\leq (\tau-\tau_1)}(\mathcal{R}^{\tau_1}(\xi_0)) \subseteq \overline{\mathcal{R}^{\leq \tau_1}(\xi_0)} \cup \overline{\mathcal{R}^{\leq (\tau-\tau_1)}(\overline{\mathcal{R}^{\tau_1}(\xi_0)})}.$$

On the other hand, combining Lemma 5.3 with the fact that $\overline{\mathcal{R}^{\leq t}(\mathcal{R}^{\tau_1}(\xi_0))}$ is compact for any $0 \leq t < \tau - \tau_1$, one sees that $\overline{\mathcal{R}^{\leq t}(\eta_1)}$ is compact for any $0 \leq t < \tau - \tau_1$.

Since $\eta_1 \in \overline{\mathcal{R}^{\tau_1}(\xi_0)}$, there exists a sequence $\{z_n\} \rightarrow \eta_1$ with $z_n \in \mathcal{R}^{\tau_1}(\xi_0)$. Assume, for each n , that $z_n = x(\tau_1, \xi_0, d_n)$ for some $d_n \in \mathcal{M}_{\mathcal{D}}$. For each $d \in \mathcal{M}_{\mathcal{D}}$, and each $s \in \mathbb{R}$, we use d_s to denote the function defined by $d_s(t) = d(s+t)$. Then by uniqueness, one has that for each n , $x(s, z_n, (d_n)_{\tau_1}) \in K_1$ for any $-\tau_1 \leq s \leq 0$, where $K_1 = \overline{\mathcal{R}^{\leq \tau_1}(\xi_0)}$. We want to claim next that, by compactness of K_1 and Gronwall's Lemma,

$$|x(-\tau_1, \eta_1, (d_n)_{\tau_1}) - \xi_0| = |x(-\tau_1, \eta_1, (d_n)_{\tau_1}) - x(-\tau_1, z_n, (d_n)_{\tau_1})| \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

The only potential problem is that the solution $x(-\tau_1, \eta_1, (d_n)_{\tau_1})$ may fail to exist a priori. However, it is possible to modify $f(x, \mathbf{d})$ outside a neighborhood of $K_1 \times \mathcal{D}$ so that it now has compact support and is hence globally bounded. The modified dynamics is complete. Now the above limit holds for the modified system, and *a fortiori* it also holds for the original system.

Choose n_0 such that

$$(28) \quad |x(-\tau_1, \eta_1, (d_{n_0})_{\tau_1}) - \xi_0| < 1/2.$$

Let $v_1 = d_{n_0}$, and let $\eta_0 = x(-\tau_1, \eta_1, (d_{n_0})_{\tau_1})$. Then, by continuity on initial conditions, there is a neighborhood \mathcal{U}_1 of η_1 contained in $B(\eta_1, 1)$ such that

$$(29) \quad |x(-\tau_1, \xi, (v_1)_{\tau_1}) - \eta_0| < 1/2, \quad \forall \xi \in \mathcal{U}_1,$$

where $B(\eta, r)$ denotes the open ball centered at η with radius r . Combining (28) and (29), one has

$$x(-\tau_1, \xi, (v_1)_{\tau_1}) \in \mathcal{U}_0, \quad \forall \xi \in \mathcal{U}_1,$$

where $\mathcal{U}_0 = B(\xi_0, 1)$.

Let $\tau_2 = \tau_1/2 = (\tau - \tau_1)/2$. Applying the above argument with ξ_0 replaced by η_1 , τ replaced by $(\tau - \tau_1)$, and τ_1 replaced by τ_2 , one shows that there exists some $\eta_2 \in \overline{\mathcal{R}^{\tau_2}(\eta_1)}$ such that $\overline{\mathcal{R}^{\leq t}(\eta_2)}$ is compact for any $0 \leq t < \tau - \sigma_2$, and $\mathcal{R}^{\leq (\tau - \sigma_2)}(\eta_2)$ is not compact, where $\sigma_2 = \tau_1 + \tau_2$, and there exist some v_2 defined on $[0, \tau_2)$ and some neighborhood \mathcal{U}_2 of η_2 contained in $B(\eta_2, 1)$, such that

$$x(-\tau_2, \xi, (v_2)_{\tau_2}) \in \mathcal{U}_1, \quad \forall \xi \in \mathcal{U}_2.$$

By induction, one can get, for each $k \geq 1$ a point η_k , a neighborhood \mathcal{U}_k of η_k contained in $B(\eta_k, 1)$, and a function v_k defined on $[0, \tau_k)$ (where $\tau_k = 2^{-k}\tau$) such that

- $\mathcal{R}^{\leq (\tau - \sigma_k)}(\eta_k)$ is not compact, where $\sigma_k = \tau_1 + \tau_2 + \dots + \tau_k = \tau(1 - 2^{-k}) \rightarrow \tau$;
- $x(-\tau_k, \xi, (v_k)_{\tau_k}) \in \mathcal{U}_{k-1}$, for any $\xi \in \mathcal{U}_k$.

Now define v on $[0, \tau)$ by concatenating all the v_k 's. That is, $v(t) = v_k(t)$ for $t \in [\sigma_{k-1}, \sigma_k)$ (with $\sigma_0 \stackrel{\text{def}}{=} 0$). Then $v \in \mathcal{M}_{\mathcal{D}}$. For each k , let

$$\zeta_k = x(-\sigma_k, \eta_k, (v^k)_{\sigma_k}),$$

where v^k is the restriction of v to $[0, \sigma_k)$. By induction,

$$x\left(-(\sigma_k - \sigma_i), \eta_k, (v^k)_{\sigma_k}\right) \in \mathcal{U}_{k-i},$$

for each $0 \leq i \leq k$, from which it follows that $\zeta_k \in \mathcal{U}_0$ for each k . By compactness of $\overline{\mathcal{U}_0}$, there exists some subsequence of $\{\zeta_k\}$ converging to some point $\zeta_0 \in \mathbb{R}^n$. For ease of notation, we still use $\{\zeta_k\}$ to denote this convergent subsequence. Our aim is next to prove that the solution starting at ζ_0 and applying the measurable function v does not exist for time τ , contradicting forward completeness.

First notice that for any compact set S , there exists some k such that $\eta_k \notin S$. Otherwise, assume that there exists some compact set S such that $\eta_k \in S$ for all k . Let $S_1 = \{\eta : d(\eta, S) \leq 1\}$. The compactness of S implies that there exists some $\delta > 0$ such that

$$\mathcal{R}^{\leq t}(\eta) \subseteq S_1$$

for any $\eta \in S$, and any $t \in [0, \delta]$. In particular, it implies that $\mathcal{R}^{\leq (\tau - \sigma_k)}(\eta_k) \subseteq S_1$ for k large enough so that $\tau - \sigma_k < \delta$. This contradicts the fact that $\mathcal{R}^{\leq (\tau - \sigma_k)}(\eta_k)$ is not compact for each k .

Assume that $x(\tau, \zeta_0, v)$ is defined. By continuity on initial conditions, this would imply that $x(t, \zeta_k, v)$ is defined for all $t \leq \tau$ and for all k large enough, and it converges uniformly to $x(t, \zeta_0, v)$. Thus, $x(t, \zeta_k, v)$ remains in a compact for all $t \in [0, \tau]$ and all k . But

$$x(\sigma_k, \zeta_k, v) = x(\sigma_k, \zeta_k, v^k) = \eta_k,$$

contradicting what was just proved. So $x(\tau, \zeta_0, v)$ is not defined, which contradicts the forward completeness of the system. \blacksquare

REMARK 5.4. For $T > 0$ and $\xi \in \mathbb{R}^n$, let

$$\mathcal{R}^{-T}(\xi) = \{\eta : \eta = x(-T, \xi, d), d \in \mathcal{M}_{\mathcal{D}}\}, \quad \text{and} \quad \mathcal{R}^{\geq -T}(\xi) = \bigcup_{t \in [-T, 0]} \mathcal{R}^t(\xi).$$

These are the reachable sets from ξ for the time reversed system

$$(30) \quad \dot{x}(t) = -f(x(t), d(t)).$$

Similarly, one defines $\mathcal{R}^{-T}(S)$ and $\mathcal{R}^{\geq -T}(S)$ for subsets S of \mathbb{R}^n . If (1) is backward complete, that is, if (30) is forward complete, and applying Proposition 5.1 to (30), one concludes, for system (1), that $\overline{\mathcal{R}^{\geq -T}(K)}$ is compact for any $T > 0$ and any compact subset K of \mathbb{R}^n . In particular, for systems that are (forward and backward) complete,

$$\overline{\mathcal{R}^{\geq -T}(K) \bigcup \mathcal{R}^{\leq T}(K)}$$

is compact for any compact set K and any $T > 0$. \square

Combining the above conclusion and Gronwall's Lemma, one has the following fact:

PROPOSITION 5.5. *Assume that (1) is complete. For any fixed $T > 0$ and any compact $K \subseteq \mathbb{R}^n$, there is a constant $C > 0$ (which only depends on the set K and T), such that for the trajectories $x(t, x_0, d)$ of the system (1),*

$$|x(t, \xi, d) - x(t, \eta, d)| \leq C|\xi - \eta|$$

for any $\xi, \eta \in K$, any $|t| \leq T$, and any $d \in \mathcal{M}_{\mathcal{D}}$. \square

6. Proof of the First Converse Lyapunov Theorem.

Proof. [\Leftarrow] Pick any $x_0 \in \mathbb{R}^n$ and any $d \in \mathcal{M}_{\mathcal{D}}$, and let $x(\cdot)$ be the corresponding trajectory. Then we have

$$\frac{dV(x(t))}{dt} \leq -\alpha_3(|x(t)|_{\mathcal{A}}) \leq -\alpha(V(x(t))), \quad \text{a.e. } t \geq 0,$$

where α is the \mathcal{K}_{∞} -function defined by

$$\alpha(\cdot) \stackrel{\text{def}}{=} \alpha_3(\alpha_2^{-1}(\cdot)).$$

Now let β_{α} be the \mathcal{KL} -function as in Lemma 4.4 with respect to α , and define

$$(31) \quad \beta(s, t) \stackrel{\text{def}}{=} \alpha_1^{-1}(\beta_{\alpha}(\alpha_2(s), t)).$$

Then β is a \mathcal{KL} -function, since both α_1 and α_2 are \mathcal{K}_{∞} -functions. By Lemma 4.4,

$$V(x(t)) \leq \beta_{\alpha}(V(x_0), t), \quad \text{any } t \geq 0.$$

Hence

$$|x(t)|_{\mathcal{A}} \leq \beta(|x_0|_{\mathcal{A}}, t), \quad \text{any } t \geq 0.$$

Therefore the system (1) is UGAS with respect to \mathcal{A} , by Proposition 2.5.

[\implies] We will show the existence of a not necessarily smooth Lyapunov function; then the existence of a smooth function will follow from Proposition 4.2. Assume that the system is UGAS with respect to the set \mathcal{A} . Let δ and T_r be as in Definition 2.2 and Lemma 3.1.

Define $g : \mathbb{R}^n \longrightarrow \mathbb{R}$ by

$$(32) \quad g(\xi) \stackrel{\text{def}}{=} \inf_{t \leq 0, d \in \mathcal{M}_{\mathcal{D}}} \{ |x(t, \xi, d)|_{\mathcal{A}} \} .$$

Note that, by uniqueness of solutions, for each $t_0 > 0$ and each d , it holds that

$$x(t - t_0, x(t_0, \xi, d), d_{t_0}) = x(t, \xi, d),$$

where d_{t_0} is defined by $d_{t_0}(t) = d(t + t_0)$. Pick any $d \in \mathcal{M}_{\mathcal{D}}$, $\xi \in \mathbb{R}^n$, and $t_1 > 0$. Let $\xi_1 = x(t_1, \xi, d)$. Then for any $t < 0$, and $v \in \mathcal{M}_{\mathcal{D}}$,

$$x(t, \xi, v) = x(t - t_1, \xi_1, v_{t_1} \# d_{t_1}),$$

where

$$v_{t_1} \# d_{t_1}(s) = \begin{cases} d(s + t_1), & \text{if } -t_1 \leq s \leq 0, \\ v(s + t_1), & \text{if } s < -t_1. \end{cases}$$

Thus,

$$\begin{aligned} g(\xi) &= \inf_{t \leq 0, v \in \mathcal{M}_{\mathcal{D}}} |x(t, \xi, v)|_{\mathcal{A}} = \inf_{t \leq 0, d \in \mathcal{M}_{\mathcal{D}}} |x(t - t_1, \xi_1, v_{t_1} \# d_{t_1})|_{\mathcal{A}} \\ &= \inf_{\tau \leq -t_1, v \in \mathcal{M}_{\mathcal{D}}} |x(\tau, \xi_1, v_{t_1} \# d_{t_1})|_{\mathcal{A}} \geq \inf_{\tau \leq 0, v \in \mathcal{M}_{\mathcal{D}}} |x(\tau, \xi_1, v)|_{\mathcal{A}} \\ &= g(\xi_1). \end{aligned}$$

This implies that

$$(33) \quad g(x(t, \xi, d)) \leq g(\xi), \quad \forall t > 0, \quad \forall d \in \mathcal{M}_{\mathcal{D}} .$$

Also one has

$$(34) \quad \delta(|\xi|_{\mathcal{A}}) \leq g(\xi) \leq |\xi|_{\mathcal{A}} .$$

The second half of (34) is obvious from $x(0, \xi, d) = \xi$. On the other hand, if the first half were not true, then there would be some $d \in \mathcal{M}_{\mathcal{D}}$ and some $t_0 \leq 0$ such that

$$\delta(|\xi|_{\mathcal{A}}) > |x(t_0, \xi, d)|_{\mathcal{A}} .$$

Pick any $0 < \varepsilon < |\xi|_{\mathcal{A}}$ so that $|x(t_0, \xi, d)|_{\mathcal{A}} < \delta(\varepsilon)$. By the uniform stability property, applied with $t = -t_0$ and $x_0 = x(t_0, \xi, d)$,

$$|\xi|_{\mathcal{A}} = |x(-t_0, x(t_0, \xi, d), d_{t_0})|_{\mathcal{A}} < |\xi|_{\mathcal{A}} ,$$

which is a contradiction.

For any $0 < \varepsilon < r$, define $K_{\varepsilon, r} \stackrel{\text{def}}{=} \{ \xi \in \mathbb{R}^n : \varepsilon \leq |\xi|_{\mathcal{A}} < r \}$.

Fact 1: For all ε and r with $0 < \varepsilon < r$, there exists $q_{\varepsilon, r} \leq 0$, such that:

$$\xi \in K_{\varepsilon, r}, \quad d \in \mathcal{M}_{\mathcal{D}}, \quad \text{and} \quad t < q_{\varepsilon, r} \implies |x(t, \xi, d)|_{\mathcal{A}} \geq r .$$

Proof: If the statement were not true, then there would exist ε, r with $0 < \varepsilon < r$ and three sequences $\{\xi_k\} \subseteq K_{\varepsilon, r}$, $\{t_k\} \subseteq \mathbb{R}$ and $d_k \in \mathcal{M}_{\mathcal{D}}$ with $\lim_{k \rightarrow \infty} t_k = -\infty$ such that for all k :

$$|x(t_k, \xi_k, d_k)|_{\mathcal{A}} < r.$$

Pick k large enough so that $-t_k > T_r(\varepsilon)$, then by the uniform attraction property,

$$|\xi_k|_{\mathcal{A}} = |x(-t_k, x(t_k, \xi_k, d_k), (d_k)_{t_k})|_{\mathcal{A}} < \varepsilon,$$

which is a contradiction. This proves the fact.

Therefore, for any $\xi \in K_{\varepsilon, r}$,

$$g(\xi) = \inf\{|x(t, \xi, d)|_{\mathcal{A}} : t \in [q_{\varepsilon, r}, 0], d \in \mathcal{M}_{\mathcal{D}}\}.$$

LEMMA 6.1. *The function $g(\xi)$ is locally Lipschitz on $\mathbb{R}^n \setminus \mathcal{A}$, and continuous everywhere.*

Proof. Fix any $\xi_0 \in \mathbb{R}^n \setminus \mathcal{A}$, and let $s = \frac{|\xi_0|_{\mathcal{A}}}{2}$. Let $\bar{B}(\xi_0, s)$ denote the closed ball centered at ξ_0 and with radius s . Then $\bar{B}(\xi_0, s) \subseteq K_{\sigma, r}$ for some $0 < \sigma < r$. Pick a constant C as in Proposition 5.5 with respect to this closed ball and $T = |q_{\sigma, r}|$. Pick any $\zeta, \eta \in \bar{B}(\xi_0, s)$. For any $\varepsilon > 0$, there exist some $d_{\eta, \varepsilon}$ and $t_{\eta, \varepsilon} \in [q_{\sigma, r}, 0]$ such that $g(\eta) \geq |x(t_{\eta, \varepsilon}, \eta, d_{\eta, \varepsilon})|_{\mathcal{A}} - \varepsilon$. Thus

$$(35) \quad g(\zeta) - g(\eta) \leq |x(t_{\eta, \varepsilon}, \zeta, d_{\eta, \varepsilon})|_{\mathcal{A}} - |x(t_{\eta, \varepsilon}, \eta, d_{\eta, \varepsilon})|_{\mathcal{A}} + \varepsilon \leq C|\zeta - \eta| + \varepsilon.$$

Note that (35) holds for all $\varepsilon > 0$, so it follows that

$$g(\zeta) - g(\eta) \leq C|\zeta - \eta|.$$

Similarly, $g(\eta) - g(\zeta) \leq C|\zeta - \eta|$. This proves that g is locally Lipschitz on $\mathbb{R}^n \setminus \mathcal{A}$.

Note that g is 0 on \mathcal{A} , and for $\xi \in \mathcal{A}$, $\eta \in \mathbb{R}^n$:

$$|g(\eta) - g(\xi)| = |g(\eta)| \leq |\eta|_{\mathcal{A}} \leq |\eta - \xi|,$$

thus g is globally continuous. (We are not claiming that g is locally Lipschitz on \mathbb{R}^n , though.) \square

Now define $U : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ by

$$(36) \quad U(\xi) \stackrel{\text{def}}{=} \sup_{t \geq 0, d \in \mathcal{M}_{\mathcal{D}}} \left\{ g(x(t, \xi, d)) k(t) \right\},$$

where $k : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{> 0}$ is any strictly increasing, smooth function that satisfies:

- there are two constants $0 < c_1 < c_2 < \infty$, such that $k(t) \in [c_1, c_2]$ for all $t \geq 0$;
- there is a bounded positive decreasing continuous function $\tau(\cdot)$, such that

$$k'(t) \geq \tau(t) \quad \text{for all } t \geq 0.$$

(For instance, $\frac{c_1 + c_2 t}{1 + t}$ is one example of such a function.) Observe that

$$(37) \quad U(\xi) \leq \sup_{t \geq 0} \left(g(\xi) k(t) \right) \leq c_2 g(\xi) \leq c_2 |\xi|_{\mathcal{A}},$$

and

$$(38) \quad U(\xi) \geq \sup_{d \in \mathcal{M}_{\mathcal{D}}} g(x(t, \xi, d)) k(t) \Big|_{t=0} \geq c_1 g(\xi) \geq c_1 \delta(|\xi|_{\mathcal{A}}).$$

For any $\xi \in \mathbb{R}^n$, since

$$|x(t, \xi, d)|_{\mathcal{A}} \leq \beta(|\xi|_{\mathcal{A}}, t), \quad \forall d, \quad \forall t \geq 0,$$

for some \mathcal{KL} -function β , and $0 \leq g(x(t, \xi, d)) \leq |x(t, \xi, d)|_{\mathcal{A}}$ for all $t \geq 0$, it follows that

$$\lim_{t \rightarrow +\infty} \sup_d g(x(t, \xi, d)) = 0.$$

Thus there exists some $\tau_\xi \in [0, \infty)$ such that

$$U(\xi) = \sup_{0 \leq t \leq \tau_\xi, d \in \mathcal{M}_{\mathcal{D}}} g(x(t, \xi, d)) k(t).$$

In fact, we can get the following explicit bound.

Fact 2: For any $0 < |\xi|_{\mathcal{A}} < r$,

$$U(\xi) = \sup_{0 \leq t \leq t_\xi, d \in \mathcal{M}_{\mathcal{D}}} g(x(t, \xi, d)) k(t),$$

where $t_\xi = T_r \left(\frac{c_1}{2c_2} \delta(|\xi|_{\mathcal{A}}) \right)$.

Proof: If the statement is not true, then for any $\varepsilon > 0$, there exists some $t_\varepsilon > T_r \left(\frac{c_1}{2c_2} \delta(|\xi|_{\mathcal{A}}) \right)$ and some d_ε such that

$$U(\xi) \leq g(x(t_\varepsilon, \xi, d_\varepsilon)) k(t_\varepsilon) + \varepsilon.$$

So we have

$$\begin{aligned} \delta(|\xi|_{\mathcal{A}}) &\leq \frac{1}{c_1} U(\xi) \leq \frac{1}{c_1} g(x(t_\varepsilon, \xi, d_\varepsilon)) k(t_\varepsilon) + \frac{\varepsilon}{c_1} \\ &\leq \frac{c_2}{c_1} g(x(t_\varepsilon, \xi, d_\varepsilon)) + \frac{\varepsilon}{c_1} \leq \frac{c_2}{c_1} |x(t_\varepsilon, \xi, d_\varepsilon)|_{\mathcal{A}} + \frac{\varepsilon}{c_1} < \frac{\delta(|\xi|_{\mathcal{A}})}{2} + \frac{\varepsilon}{c_1}. \end{aligned}$$

Taking the limit as ε tends to 0 results in a contradiction.

For any compact set $K \subseteq \mathbb{R}^n \setminus \mathcal{A}$, let

$$t_K \stackrel{\text{def}}{=} \max_{\xi \in K} t_\xi < \infty.$$

(Finiteness follows from Fact 2, as $K \subseteq \{\xi : 0 < |\xi|_{\mathcal{A}} < r\}$ for some $r > 0$.)

LEMMA 6.2. *The function $U(\cdot)$ defined by (36) is locally Lipschitz on $\mathbb{R}^n \setminus \mathcal{A}$, and continuous everywhere.*

Proof. For $\xi_0 \notin \mathcal{A}$, pick up a compact neighborhood K_0 of ξ_0 so that $K_0 \cap \mathcal{A} = \emptyset$. By (38), one knows that

$$U(\xi) > r_0, \quad \forall \xi \in K_0,$$

for some constant $r_0 > 0$. Let $r_1 = \frac{r_0}{2c_2}$ and let

$$K_1 = K_0 \cap \left\{ \eta : |\eta - \xi_0| \leq \frac{r_1}{4C} \right\},$$

where C is such a constant that

$$(39) \quad |x(t, \xi, d) - x(t, \eta, d)| \leq C |\xi - \eta|, \quad \forall \xi, \eta \in K_0, \quad 0 \leq t \leq t_{K_0}, \quad d \in \mathcal{M}_{\mathcal{D}}.$$

In the following we will show that there exists some $L > 0$ such that for any $\xi, \eta \in K_1$, it holds that

$$(40) \quad |U(\xi) - U(\eta)| \leq L |\xi - \eta|.$$

First of all, for any $\xi \in K_1$ and any $\varepsilon \in (0, r_0/2)$, there exists $t_{\xi,\varepsilon} \in [0, t_{K_0}]$ and $d_{\xi,\varepsilon} \in \mathcal{M}_{\mathcal{D}}$, such that

$$U(\xi) \leq g(x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon}))k(t_{\xi,\varepsilon}) + \varepsilon \leq c_2 |x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon})|_{\mathcal{A}} + \varepsilon,$$

from which it follows that

$$|x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon})|_{\mathcal{A}} \geq r_1.$$

It follows from (39) that for any $\eta \in K_1$,

$$|x(t_{\xi,\varepsilon}, \eta, d_{\xi,\varepsilon})|_{\mathcal{A}} \geq |x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon})|_{\mathcal{A}} - |x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon}) - x(t_{\xi,\varepsilon}, \eta, d_{\xi,\varepsilon})| \geq \frac{r_1}{2}.$$

By Proposition 5.1 one knows that there exists some compact set K_2 such that

$$x(t, \xi, d) \in K_2, \quad \forall \xi \in K_1, \quad \forall t \in [0, t_{K_1}], \quad \text{and } \forall d \in \mathcal{M}_{\mathcal{D}}.$$

Again, applying Lemma 6.1 to the compact set $K_2 \cap \{\zeta : |\zeta|_{\mathcal{A}} \geq r_1/2\}$, one sees that

$$|g(x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon})) - g(x(t_{\xi,\varepsilon}, \eta, d_{\xi,\varepsilon}))| \leq C_1 |x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon}) - x(t_{\xi,\varepsilon}, \eta, d_{\xi,\varepsilon})|,$$

for some $C_1 > 0$. Therefore, we have the following:

$$\begin{aligned} U(\xi) - U(\eta) &\leq g(x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon}))k(t_{\xi,\varepsilon}) + \varepsilon - g(x(t_{\xi,\varepsilon}, \eta, d_{\xi,\varepsilon}))k(t_{\xi,\varepsilon}) \\ &\leq c_2 |g(x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon})) - g(x(t_{\xi,\varepsilon}, \eta, d_{\xi,\varepsilon}))| + \varepsilon \\ &\leq C_1 c_2 |x(t_{\xi,\varepsilon}, \xi, d_{\xi,\varepsilon}) - x(t_{\xi,\varepsilon}, \eta, d_{\xi,\varepsilon})| + \varepsilon \\ &\leq L |\xi - \eta| + \varepsilon, \end{aligned}$$

for some constant L that only depends on the compact set K_1 . Note that the above holds for any $\varepsilon \in (0, r_0/2)$, thus,

$$U(\xi) - U(\eta) \leq L |\xi - \eta|, \quad \forall \xi, \eta \in K_1.$$

By symmetry, one proves (40).

To prove the continuity of U on \mathbb{R}^n , note that for any $\xi \in \mathcal{A}$, it holds that $U(\xi) = 0$, and so for all $\eta \in \mathbb{R}^n$:

$$|U(\xi) - U(\eta)| = U(\eta) \leq c_2 |\eta|_{\mathcal{A}} \leq c_2 |\xi - \eta|.$$

The proof of Lemma 6.2 is thus concluded. \square

We next start proving that U decreases along trajectories. Now pick any $\xi \notin \mathcal{A}$. Let $h_0 > 0$ be such that

$$|x(t, \xi, \mathbf{d})|_{\mathcal{A}} \geq \frac{|\xi|_{\mathcal{A}}}{2}, \quad \forall \mathbf{d} \in \mathcal{D}, \quad \forall t \in [0, h_0],$$

where \mathbf{d} denotes the constant function $d(t) \equiv \mathbf{d}$. Such an h_0 exists by continuity. Pick any $h \in [0, h_0]$. For each $\mathbf{d} \in \mathcal{D}$, let $\eta_{\mathbf{d}} = x(h, \xi, \mathbf{d})$. For any $\varepsilon > 0$, there exist some $t_{\mathbf{d},\varepsilon}$ and $d_{\mathbf{d},\varepsilon} \in \mathcal{M}_{\mathcal{D}}$ such that

$$\begin{aligned} U(\eta_{\mathbf{d}}) &\leq g(x(t_{\mathbf{d},\varepsilon}, \eta_{\mathbf{d}}, d_{\mathbf{d},\varepsilon}))k(t_{\mathbf{d},\varepsilon}) + \varepsilon \\ &= g(x(t_{\mathbf{d},\varepsilon} + h, \xi, \tilde{d}_{\mathbf{d},\varepsilon}))k(t_{\mathbf{d},\varepsilon} + h) \left(1 - \frac{k(t_{\mathbf{d},\varepsilon} + h) - k(t_{\mathbf{d},\varepsilon})}{k(t_{\mathbf{d},\varepsilon} + h)}\right) + \varepsilon \\ (41) \quad &\leq U(\xi) \left(1 - \frac{k(t_{\mathbf{d},\varepsilon} + h) - k(t_{\mathbf{d},\varepsilon})}{c_2}\right) + \varepsilon, \end{aligned}$$

where $\tilde{\mathbf{d}}_{\mathbf{d},\varepsilon}$ is the concatenation of \mathbf{d} and $d_{\mathbf{d},\varepsilon}$. Still for these ξ and h , and for any $r > |\xi|_{\mathcal{A}}$, define

$$(42) \quad T_{\xi,h}^r \stackrel{\text{def}}{=} \max_{0 \leq \bar{t} \leq h, \mathbf{d} \in \mathcal{D}} T_r \left(\frac{c_1}{2c_2} \delta(|x(\bar{t}, \xi, \mathbf{d})|_{\mathcal{A}}) \right).$$

Claim: $t_{\mathbf{d},\varepsilon} + h \leq T_{\xi,h}^r$, for all $\mathbf{d} \in \mathcal{D}$ and for all $\varepsilon \in \left(0, \frac{c_1}{2} \delta \left(\frac{|\xi|_{\mathcal{A}}}{2} \right)\right)$.

Proof: If this were not true, then there would exist some $\tilde{\mathbf{d}}$ and some $\tilde{\varepsilon} \in \left(0, \frac{c_1}{2} \delta \left(\frac{|\xi|_{\mathcal{A}}}{2} \right)\right)$ such that $t_{\tilde{\mathbf{d}},\tilde{\varepsilon}} + h > T_{\xi,h}^r$, and hence in particular, for $\bar{t} = h$ and $\mathbf{d} = \tilde{\mathbf{d}}$ it holds that

$$t_{\tilde{\mathbf{d}},\tilde{\varepsilon}} + h > T_r \left(\frac{c_1}{2c_2} \delta(|\eta_{\tilde{\mathbf{d}}}|_{\mathcal{A}}) \right),$$

which implies that

$$|x(t_{\tilde{\mathbf{d}},\tilde{\varepsilon}}, \eta_{\tilde{\mathbf{d}}}, d_{\tilde{\mathbf{d}},\tilde{\varepsilon}})|_{\mathcal{A}} = |x(t_{\tilde{\mathbf{d}},\tilde{\varepsilon}} + h, \xi, v)|_{\mathcal{A}} < \frac{c_1}{2c_2} \delta(|\eta_{\tilde{\mathbf{d}}}|_{\mathcal{A}}),$$

where v is the concatenated function defined by

$$v(t) = \begin{cases} \tilde{\mathbf{d}}, & \text{if } 0 \leq t \leq h, \\ d_{\tilde{\mathbf{d}},\tilde{\varepsilon}}(t-h), & \text{if } t > h. \end{cases}$$

Using (38), one has

$$\begin{aligned} \delta(|\eta_{\tilde{\mathbf{d}}}|_{\mathcal{A}}) &\leq \frac{1}{c_1} U(\eta_{\tilde{\mathbf{d}}}) \leq \frac{1}{c_1} g(x(t_{\tilde{\mathbf{d}},\tilde{\varepsilon}}, \eta_{\tilde{\mathbf{d}}}, d_{\tilde{\mathbf{d}},\tilde{\varepsilon}})) k(t_{\tilde{\mathbf{d}},\tilde{\varepsilon}}) + \frac{\tilde{\varepsilon}}{c_1} \\ &\leq \frac{c_2}{c_1} |x(t_{\tilde{\mathbf{d}},\tilde{\varepsilon}}, \eta_{\tilde{\mathbf{d}}}, d_{\tilde{\mathbf{d}},\tilde{\varepsilon}})|_{\mathcal{A}} + \frac{\tilde{\varepsilon}}{c_1} < \frac{1}{2} \delta(|\eta_{\tilde{\mathbf{d}}}|_{\mathcal{A}}) + \frac{\tilde{\varepsilon}}{c_1}, \end{aligned}$$

which is a contradiction, since $\tilde{\varepsilon} < \frac{c_1}{2} \delta \left(\frac{|\xi|_{\mathcal{A}}}{2} \right) \leq \frac{c_1 \delta(|\eta_{\tilde{\mathbf{d}}}|_{\mathcal{A}})}{2}$. This proves the claim.

From (41), we have for any $\mathbf{d} \in \mathcal{D}$ and for any $\varepsilon > 0$ small enough,

$$U(x(h, \xi, \mathbf{d})) - U(\xi) \leq -U(\xi) \frac{(k(t_{\mathbf{d},\varepsilon} + h) - k(t_{\mathbf{d},\varepsilon}))}{c_2} + \varepsilon = -\frac{U(\xi)}{c_2} k'(t_{\mathbf{d},\varepsilon} + \theta h) h + \varepsilon,$$

where θ is some number in $(0, 1)$. Hence, by the assumptions made on the function k , we have

$$U(x(h, \xi, \mathbf{d})) - U(\xi) \leq -\frac{U(\xi)}{c_2} \tau(t_{\mathbf{d},\varepsilon} + \theta h) h + \varepsilon \leq -\frac{U(\xi)}{c_2} \tau(T_{\xi,h}^r) h + \varepsilon.$$

Again, since ε can be chosen arbitrarily small, we have

$$U(x(h, \xi, \mathbf{d})) - U(\xi) \leq -\frac{U(\xi)}{c_2} \tau(T_{\xi,h}^r) h, \quad \forall \mathbf{d} \in \mathcal{D}.$$

Thus we showed that for any \mathbf{d} and any $h > 0$ small enough,

$$\frac{U(x(h, \xi, \mathbf{d})) - U(\xi)}{h} \leq -\frac{U(\xi)}{c_2} \tau(T_{\xi,h}^r).$$

Since U is locally Lipschitz on $\mathbb{R}^n \setminus \mathcal{A}$, it is differentiable almost everywhere in $\mathbb{R}^n \setminus \mathcal{A}$, and hence for any $\mathbf{d} \in \mathcal{D}$ and for any $r > |\xi|_{\mathcal{A}}$,

$$\begin{aligned} L_{f_{\mathbf{d}}} U(\xi) &= \lim_{h \rightarrow 0^+} \frac{U(x(h, \xi, \mathbf{d})) - U(\xi)}{h} \leq -\lim_{h \rightarrow 0^+} \frac{U(\xi)}{c_2} \tau(T_{\xi,h}^r) \\ &= -\frac{U(\xi)}{c_2} \tau \left(\lim_{h \rightarrow 0^+} T_{\xi,h}^r \right) = -\frac{U(\xi)}{c_2} \tau \left(T_r \left(\frac{c_1}{2c_2} \delta(|\xi|_{\mathcal{A}}) \right) \right) \\ (43) \quad &\leq -\frac{c_1 \delta(|\xi|_{\mathcal{A}})}{c_2} \tau \left(T_r \left(\frac{c_1}{2c_2} \delta(|\xi|_{\mathcal{A}}) \right) \right) \\ (44) \quad &= -\bar{\alpha}_r(|\xi|_{\mathcal{A}}), \text{ a.e. ,} \end{aligned}$$

where

$$\bar{\alpha}_r(s) = \frac{c_1 \delta(s)}{c_2} \tau \left(T_r \left(\frac{c_1}{2c_2} \delta(s) \right) \right).$$

Now define the function $\bar{\alpha}$ by

$$\bar{\alpha}(s) = \sup_{r>s} \bar{\alpha}_r(s).$$

Note that $\bar{\alpha}_r(0) = 0$ for any $r > 0$, so $\bar{\alpha}(0) = 0$. Also, applying to $r = 2s$, we have

$$\bar{\alpha}(s) \geq \frac{c_1 \delta(s)}{c_2} \tau \left(T_{2s} \left(\frac{c_1}{2c_2} \delta(s) \right) \right) > 0$$

for all $s > 0$. Notice that (44) holds for any $r > |\xi|_{\mathcal{A}}$, so it follows that for every $\mathbf{d} \in \mathcal{D}$, $L_{f_{\mathbf{d}}}U(\xi) \leq -\bar{\alpha}(|\xi|_{\mathcal{A}})$ for almost all $\xi \in \mathbb{R}^n \setminus \mathcal{A}$. Now let

$$\check{\alpha}(s) = \frac{c_1 \delta(s)}{c_2} \int_{2s}^{2s+1} \tau \left(T_r \left(\frac{c_1}{2c_2} \delta(s) \right) \right) dr,$$

for $s > 0$, and let $\check{\alpha}(0) = 0$. Then $\check{\alpha}$ is continuous on $[0, \infty)$ (the continuity at $s = 0$ is because τ is bounded and $\delta(0) = 0$), and for $s > 0$, it holds that

$$0 < \check{\alpha}(s) \leq \frac{c_1 \delta(s)}{c_2} \tau \left(T_{2s} \left(\frac{c_1}{2c_2} \delta(s) \right) \right)$$

because of the monotonicity properties of T and τ . Furthermore,

$$L_{f_{\mathbf{d}}}U(\xi) \leq -\bar{\alpha}(|\xi|_{\mathcal{A}}) \leq -\check{\alpha}(|\xi|_{\mathcal{A}}),$$

for almost all $\xi \in \mathbb{R}^n \setminus \mathcal{A}$.

By Theorem B.1 provided in the appendix, there exists a C^∞ function $V : \mathbb{R}^n \setminus \mathcal{A} \rightarrow \mathbb{R}_{\geq 0}$ such that for almost all $\xi \in \mathbb{R}^n \setminus \mathcal{A}$,

$$|V(\xi) - U(\xi)| < \frac{U(\xi)}{2} \quad \text{and} \quad L_{f_{\mathbf{d}}}V(\xi) \leq -\frac{1}{2}\check{\alpha}(|\xi|_{\mathcal{A}}), \quad \forall \mathbf{d} \in \mathcal{D}.$$

Extend V to \mathbb{R}^n by letting $V|_{\mathcal{A}} = 0$ and again denote the extension by V . Note that V is continuous on \mathbb{R}^n . So V is a Lyapunov function, as desired, with $\alpha_1(s) = \frac{c_1}{2}\delta(s)$, $\alpha_2(s) = \frac{3c_2}{2}s$ and $\alpha_3(s) = \frac{1}{2}\check{\alpha}(s)$. \square

7. Proof of the Second Converse Lyapunov Theorem. We need a couple of Lemmas. The first one is trivial, so we omit its proof.

LEMMA 7.1. *Let $f : \mathbb{R}^n \times \mathcal{D} \rightarrow \mathbb{R}^n$ be continuous, where \mathcal{D} is a compact subset of \mathbb{R}^l . Then there exists a smooth function $a_f : \mathbb{R}^n \rightarrow \mathbb{R}$, with $a_f(x) \geq 1$ everywhere, such that $|f(x, \mathbf{d})| \leq a_f(x)$ for all x and all \mathbf{d} .* \square

Now for any given system

$$\Sigma : \dot{x} = f(x, \mathbf{d}),$$

not necessarily complete, consider the following system:

$$\Sigma_b : \dot{x} = \frac{1}{a_f(x)} f(x, \mathbf{d}).$$

Note that the system Σ_b is complete since $\frac{|f(x, \mathbf{d})|}{a_f(x)} \leq 1$ for all x, \mathbf{d} . We let $x_b(\cdot, x_0, d)$ denote the trajectory of Σ_b corresponding to the initial state x_0 and the time-varying parameter d . The following result is a simple

consequence of the fact that the trajectories of Σ are the same as those of Σ_b up to a rescaling of time. We provide the details to show clearly that the uniformity conditions are not violated.

LEMMA 7.2. *Assume that \mathcal{A} is a compact set. Suppose that system Σ is UGAS with respect to \mathcal{A} . Then, system Σ_b is UGAS with respect to \mathcal{A} .*

Proof. Pick a time-varying parameter $d \in \mathcal{M}_{\mathcal{D}}$ and an initial state $x_0 \in \mathbb{R}^n$. Let $\gamma_b(t)$ denote $x_b(t, x_0, d)$. Let $\tau_{\gamma_b}(t)$ denote the solution for $t \geq 0$ of the following initial value problem:

$$(45) \quad \dot{\tau} = a_f(\gamma_b(\tau)), \quad \tau(0) = 0.$$

Since a_f is smooth, and γ_b is Lipschitz, $a_f \circ \gamma_b$ is locally Lipschitz as well. It follows that a unique $\tau_{\gamma_b}(t)$ is at least defined in some interval $[0, \bar{t})$. Note that τ_{γ_b} is strictly increasing, so $\bar{t} < +\infty$ would imply $\lim_{t \rightarrow \bar{t}^-} \tau_{\gamma_b}(t) = +\infty$.

Claim: For every trajectory γ_b of Σ_b , $\tau_{\gamma_b}(t)$ is defined for all $t \geq 0$.

Proof: If the claim is not true, then there exist some trajectory γ_b of Σ_b and some $t_1 > 0$ such that $\lim_{t \rightarrow t_1^-} \tau_{\gamma_b}(t) = \infty$. Now for $t \in [0, t_1)$, one has:

$$(46) \quad \begin{aligned} \frac{d}{dt} \gamma_b(\tau_{\gamma_b}(t)) &= \frac{1}{a_f(\gamma_b(\tau_{\gamma_b}(t)))} f\left(\gamma_b(\tau_{\gamma_b}(t)), d(\tau_{\gamma_b}(t))\right) \frac{d}{dt} \tau_{\gamma_b}(t) \\ &= f\left(\gamma_b(\tau_{\gamma_b}(t)), d(\tau_{\gamma_b}(t))\right). \end{aligned}$$

Thus $\gamma_b(\tau_{\gamma_b}(t))$ is a solution of Σ on $[0, t_1)$. By the stability of Σ , it follows that

$$|\gamma_b(\tau_{\gamma_b}(t))|_{\mathcal{A}} < \delta^{-1}(|x_0|_{\mathcal{A}}), \quad t \in [0, t_1),$$

where $x_0 = \gamma_b(0)$, and δ is the function for Σ as defined in Definition 2.2. (c.f. Remark 2.4.) Let $c = \delta^{-1}(|x_0|_{\mathcal{A}})$, and let $M = \sup_{|\xi|_{\mathcal{A}} \leq c} a_f(\xi)$. (M is finite because the set $\{\xi : |\xi|_{\mathcal{A}} \leq c\}$ is a compact set.) From here one sees that $|\tau_{\gamma_b}(t)| \leq Mt_1$ for any $t \in [0, t_1)$. This is a contradiction. Thus $\tau_{\gamma_b}(t)$ is defined for all $t \geq 0$. This proves the claim.

Since $a_f(s) \geq 1$ and, for every trajectory γ_b of Σ_b , $\tau_{\gamma_b}(0) = 0$, it follows that $\tau_{\gamma_b}(\cdot) \in \mathcal{K}_{\infty}$ for each trajectory γ_b of Σ_b . From (46), one also sees that if $\gamma_b(t)$ is a trajectory of Σ_b , then $\gamma_b(\tau_{\gamma_b}(t))$ is a trajectory of Σ , and furthermore,

$$|\gamma_b(\tau_{\gamma_b}(s))|_{\mathcal{A}} < \varepsilon \quad \forall s \geq 0, \quad \text{if } |\gamma_b(0)|_{\mathcal{A}} \leq \delta(\varepsilon).$$

It follows that

$$|\gamma_b(t)|_{\mathcal{A}} = |\gamma_b(\tau_{\gamma_b}(\tau_{\gamma_b}^{-1}(t)))|_{\mathcal{A}} < \varepsilon, \quad \forall t \geq 0, \quad \text{whenever } |\gamma_b(0)|_{\mathcal{A}} \leq \delta(\varepsilon).$$

This shows that condition (1) of Definition 2.2 holds for Σ_b , with the same function δ .

Fix any $r, \varepsilon > 0$. Pick any x_0 with $|x_0|_{\mathcal{A}} < r$ and any $d \in \mathcal{M}_{\mathcal{D}}$. Again let $\gamma_b(t)$ denote the corresponding trajectory of Σ_b . Then

$$|\gamma_b(t)|_{\mathcal{A}} = |\gamma_b(\tau_{\gamma_b}(\tau_{\gamma_b}^{-1}(t)))|_{\mathcal{A}} < \delta^{-1}(r), \quad \forall t \geq 0.$$

Let

$$L = \sup\{a_f(\xi) : |\xi|_{\mathcal{A}} \leq \delta^{-1}(r)\}.$$

Then one sees that $|\dot{\tau}(t)| \leq L$, which implies that $\tau_{\gamma_b}(t) \leq Lt$ for all $t \geq 0$. Note that for the given $r, \varepsilon > 0$, by the UGAS property for Σ , there exists $T > 0$ such that for every $d \in \mathcal{M}_{\mathcal{D}}$,

$$|\gamma_b(\tau_{\gamma_b}(s))|_{\mathcal{A}} < \varepsilon$$

whenever $|\gamma_b(0)|_{\mathcal{A}} < r$ and $s \geq T$. This implies that

$$|\gamma_b(t)|_{\mathcal{A}} < \varepsilon$$

whenever $|\gamma_b(0)|_{\mathcal{A}} < r$ and $t \geq \tau_{\gamma_b}(T)$. Combining this with the fact that $\tau_{\gamma_b}(t) \leq Lt$, one proves that for any $d \in \mathcal{M}_{\mathcal{D}}$, it holds that

$$|\gamma_b(t)|_{\mathcal{A}} < \varepsilon$$

whenever $|\gamma_b(0)|_{\mathcal{A}} < r$ and $t \geq LT$. Hence we conclude that Σ_b is UGAS. \square

In Lemma 7.2, the assumption that \mathcal{A} is compact is crucial. Without this assumption, the conclusion may fail as the following example shows.

EXAMPLE 7.3. Consider the following system Σ :

$$(47) \quad \dot{x} = (1 + y^2) \tanh x, \quad \dot{y} = y^4.$$

(Here f is independent of d .) Let $\mathcal{A} = \{(x, y) : x = 0\}$. Clearly the system is UGAS with respect to \mathcal{A} . For this system, a natural choice of a_f is $2 + y^4$. Thus, the corresponding Σ_b is as follows:

$$\dot{x} = (\tanh x) \frac{1 + y^2}{2 + y^4}, \quad \dot{y} = \frac{y^4}{2 + y^4}.$$

However, the system Σ_b is not UGAS with respect to \mathcal{A} . This can be seen as follows. Assume that Σ_b is UGAS. Then for $\varepsilon = 1/2$, there exists some $T > 0$ such that for any solution $(x(t), y(t))$ of Σ_b with $x(0) = 1$, it holds that

$$(48) \quad |x(t)| < \frac{1}{2}, \quad \forall t \geq T.$$

Since $\frac{1 + y^2}{2 + y^4} \rightarrow 0$ as $y \rightarrow \infty$, it follows that there exists some $y_0 > 0$ such that

$$\left| \frac{1 + y^2}{2 + y^4} \right| < \frac{1}{3T}, \quad \forall y \geq y_0.$$

Now consider the trajectory $(x(t), y(t))$ of Σ_b with $x(0) = 1, y(0) = y_0$, where y_0 is as above. Clearly $y(t) \geq y_0$ for all $t \geq 0$, and thus,

$$\dot{x} = (\tanh x) \frac{1 + y^2}{2 + y^4} \leq (\tanh x) \frac{1}{3T} \leq \frac{1}{3T},$$

which implies that

$$|x(T)| \geq 1 - \frac{1}{3T} T = \frac{2}{3}.$$

This contradicts (48). From here one sees that Σ_b is not UGAS with respect to \mathcal{A} . \square

We now prove Theorem 2.9.

The proof of the sufficiency part is the same as in the proof of Theorem 2.8. Observe that the fact that $V(\xi)$ is nonincreasing along trajectories implies, by compactness of \mathcal{A} , that trajectories are bounded, so $x(t)$ is defined for all $t \geq 0$. We now prove necessity.

Let a_f be a function for f as in Lemma 7.1, and let Σ_b be the corresponding system. Then by Lemma 7.2, one knows that the system Σ_b is UGAS. Applying Theorem 2.8 to the complete system Σ_b , one knows that there exists a smooth Lyapunov function V for Σ_b such that

$$\alpha_1(|\xi|_{\mathcal{A}}) \leq V(\xi) \leq \alpha_2(|\xi|_{\mathcal{A}}), \quad \forall \xi \in \mathbb{R}^n,$$

and

$$L_{\tilde{f}_{\mathbf{d}}} V(\xi) \leq -\alpha_3(|\xi|_{\mathcal{A}}), \quad \forall \xi \notin \mathcal{A}, \quad \forall \mathbf{d} \in \mathcal{D},$$

for some \mathcal{K}_∞ -functions α_1, α_2 and some positive definite function α_3 , where

$$\tilde{f}_{\mathbf{d}}(\xi) = \frac{f(\xi, \mathbf{d})}{a_f(\xi)}.$$

Since $a_f(\xi) \geq 1$ everywhere, it follows that

$$L_{f_{\mathbf{d}}} V(\xi) \leq -\alpha_3(|\xi|_{\mathcal{A}}), \quad \forall \xi \notin \mathcal{A}, \quad \forall \mathbf{d} \in \mathcal{D}.$$

Thus, one concludes that V is also a Lyapunov function of Σ .

8. An Example. In general, for a noncompact parameter value set \mathcal{D} , the converse Lyapunov theorem will fail, even if the vector fields $f(\xi, \mathbf{d})$ are locally Lipschitz uniformly on \mathbf{d} on any compact subset of \mathcal{D} (for instance, if f is smooth everywhere). To illustrate this fact, consider the common case of systems affine in controls:

$$\dot{x} = f(x) + g(x)\mathbf{d},$$

where for simplicity we consider only the unconstrained single-input case, that is, $\mathcal{D} = \mathbb{R}$. Assume that there would exist a Lyapunov function V for this system in the sense of definition 2.6. Then, calculating Lie derivatives, we have that, in particular,

$$L_f V(\xi) + \mathbf{d}L_g V(\xi) < 0, \quad \forall \xi \neq 0, \quad \forall \mathbf{d} \in \mathbb{R},$$

which implies that

$$L_g V(\xi) = 0, \quad \forall \xi \neq 0.$$

Thus V must be constant along all the trajectories of the differential equation

$$\dot{x} = g(x).$$

In general, such a property will contradict the properness or the positive definiteness of V , unless the vector field g is very special. As a way to construct counterexamples, consider the following property of a vector field g , which is motivated by the prolongation ideas in [28].

Consider the closure $W(\xi_0)$ of the trajectory through ξ_0 with respect to the vector field g . Note that if $\xi_1 \in W(\xi_0)$, then the fact that V is constant on trajectories, coupled with continuity of V , implies that $V(\xi_1) = V(\xi_0)$. Now assume that there is a chain $\xi_0, \xi_1, \xi_2, \dots$ so that for each $i = 1, 2, \dots$, $\xi_i \in W(\xi_{i-1})$. Then we conclude that $V(\xi_i) = V(\xi_0)$ for all i . If the sequence $\{\xi_i\}$ converges to zero (and $\xi_0 \neq 0$) or diverges to infinity, we contradict positive definiteness or properness of V respectively. For an example, take the following two dimensional system, which was used in [7] to show essentially the same fact.

Let \mathfrak{S} be the spiral that describes the solution of the differential equation

$$\dot{x} = -x - y, \quad \dot{y} = x - y,$$

passing through the point $(1, 0)$. Explicitly, \mathfrak{S} can be parameterized as $x = e^{-t} \cos t$, $y = e^{-t} \sin t$, $-\infty < t < \infty$. In polar coordinates, the spiral is given by $r = e^{-\theta}$, $-\infty < \theta < \infty$. Let $a(x, y)$ be any nonnegative smooth function which is zero exactly on the closure of the spiral \mathfrak{S} (that is, \mathfrak{S} plus the origin). (Such a function always

exists since any closed subset of Euclidean space can be described as the zero set of a smooth function; see for instance [6]. Now consider the system

$$(49) \quad \begin{aligned} \dot{x} &= -x - y + xa(x, y)\mathbf{d}, \\ \dot{y} &= x - y + ya(x, y)\mathbf{d}. \end{aligned}$$

Note that the system is smooth everywhere. Let $\mathcal{D} = \mathbb{R}$, and let \mathcal{A} be the origin. In polar coordinates, the system (49) on $\mathbb{R}^2 \setminus \{0\}$ satisfies the equations

$$(50) \quad \dot{r} = -r + ra(r \cos \theta, r \sin \theta)\mathbf{d}, \quad \dot{\theta} = 1.$$

(This can be seen as a system on $\mathbb{R}_{>0} \times S^1$.) In polar coordinates, then, the trajectory passing through $(r, \theta) = (1, 0)$ is precisely the spiral $r = e^{-\theta}$, for any $d \in \mathcal{M}_{\mathcal{D}}$. Pick any trajectory $(r(t), \theta(t))$ with $(r(0), \theta(0)) = (r_0, \theta_0)$, where $\theta_0 \in [0, 2\pi)$. Then there exists some integer $k \geq 0$ such that $r_0 < e^{-\theta_0 + 2k\pi}$.

Claim: It holds that

$$(51) \quad r(t) \leq e^{-\theta_0 + 2k\pi - t} \leq e^{2k\pi - t}, \quad \forall t \geq 0.$$

Assume that (51) is not true. Then there exists some $t_1 > 0$ such that

$$r(t_1) = e^{-\theta_0 + 2k\pi - t_1}.$$

Note that we also have $\theta(t_1) = \theta_0 + t_1$. Now let $(\bar{r}(t), \bar{\theta}(t)) = (e^{-\theta_0 + 2k\pi - t}, \theta_0 - 2k\pi + t)$. Then $(\bar{r}(t), \bar{\theta}(t))$ is a trajectory of the system, and furthermore, $(\bar{r}(0), \bar{\theta}(0))$ and $(r(0), \theta_0)$ are different points since $\bar{r}(0) \neq r(0)$. However, the points $(r(t_1), \theta(t_1))$ and $(\bar{r}(t_1), \bar{\theta}(t_1))$ are the same point on the xy -plane. This violates the uniqueness of solutions. Therefore, (51) holds for $t \geq 0$.

Note that in the above discussion, one can always choose $k = r_0 + 1$. It then follows from (51) that for any trajectory of the system with $r(0) = r_0$, it holds that

$$(52) \quad r(t) \leq e^{2(r_0+1)\pi - t}, \quad \forall t \geq 0, \quad \forall d.$$

Thus we conclude that the system is UGAS.

However, this system fails to admit a Lyapunov function. In this example, the vector field g is $(xa(x, y), ya(x, y))$. Consider the sequence of points in the xy -plane $\{\xi_k\}$ with $\xi_k = (e^{2k\pi}, 0)$ for $k \geq 0$. Note that for each $k \geq 1$,

$$\xi_k \in W(\xi_{k-1}^j),$$

where $\xi_k^j = \left(e^{2k\pi} + \frac{1}{j}, 0\right)$. Therefore, $V(\xi_k) = V(\xi_{k-1}^j)$ for any j and any k . This implies that

$$V(\xi_k) = V(\xi_0), \quad \forall k \geq 1,$$

contradicting the properness of V . This shows that it is impossible for the system to have a Lyapunov function.

It is worthwhile to note that by the same argument, one sees that not only there is no smooth Lyapunov function for the system, but also there is not even a Lyapunov function which is merely continuous (in the sense that V is not even smooth away from \mathcal{A} , and the Lie derivative condition is replaced by a condition asking that V should decrease along trajectories).

In [17], a simple example is given illustrating that uniform global asymptotic stability with respect merely to *constant* parameters is also not sufficient to guarantee the existence of Lyapunov functions.

9. Relations to Other Work. The study of smooth Converse Lyapunov Theorems has a long history. In the special case of stability with respect to equilibria, and for systems without parameters, the first complete work was that done in the early 1950s by Massera and Kurzweil; see for instance the papers [18] and [13]. (Although more general because we deal with set stability and time varying parameters, there is one important aspect in which our results are weaker than some of this classical work, especially that of Kurzweil: we assume enough regularity on the original system, so that there are unique solutions and there is continuous dependence. We do so because lack of regularity is not an issue in the main applications in which we are interested. Of course, the proofs become much simpler under regularity assumptions.) In the late 1960s, Wilson, in [31], extended the Massera and Kurzweil results to a converse Lyapunov function theorem for local asymptotic stability with respect to closed sets. Several details of critical steps were omitted in [31]. In 1990, Nadzieja in [21] filled-in the missing steps of the proof in [31], but only for the special case when the invariant set is compact. As explained earlier, our proof is also modeled along the lines [31]. See also the textbooks [32] and [12] for many of these classical results.

Nondifferentiable Lyapunov functions have been studied in many papers and textbooks. Among these we may mention the classic book [3] by Bhatia and Szegö, as well as Zubov’s work (see for instance [33]) which study in detail continuous Lyapunov function characterizations for global asymptotic stability with respect to arbitrary closed invariant sets. Also, in [29] and [28], and related work, the authors obtained the existence of continuous Lyapunov functions for systems which are stable, uniformly on parameters (or inputs) and with respect to compact sets, assuming various additional conditions involving prolongations of dynamical systems. (The next section provides some more details on the prolongation approach.) Many results on converse Lyapunov functions with respect to sets can also be found in the many books and articles by Lakshmikantham and several coauthors. For instance, in [14], Theorem 3.4.1, a Massera-type proof is provided of a general converse theorem on local asymptotic stability with respect to two \mathcal{K} -functions, that provides a Lipschitz Lyapunov function. As the authors point out, their theorem immediately provides a set-stability result (when using distance to the set as one of the comparison functions). In the very recent work [22], the author considered asymptotic stability for systems with merely measurable right hand sides, and proved the existence of locally Lipschitz Lyapunov functions for such systems. Note that in our case, we obtained the existence of locally Lipschitz Lyapunov functions as an intermediate result, but our regularity assumption on the vector fields made it possible to obtain the existence of smooth Lyapunov functions.

The questions addressed in this paper are related to studies of “total stability,” which typically ask about the preservation of stability when considering a new system $\dot{x} = f(x) + R(x, t)$, where $R(x, t)$ is a perturbation. (Sometimes the original system may be allowed to be time-varying, that is, it has equations $\dot{x} = f(x, t)$; in that case, its stability can in turn be interpreted in terms of stability of the set $\{x = 0\}$ for the extended system $\dot{x} = f(x, z)$, $\dot{z} = 1$.) In [15], Lefschetz discussed stability with respect to equilibria under perturbations (referred to by the author as quasi-stability). In [12] and [32], one can find such studies, and relationships to the special case of $\dot{x} = f(x) + d(t)$, with results proved regarding stability under integrable perturbations (not arbitrary bounded ones).

Under suitable technical conditions, systems with time varying parameters can also be treated as general dynamical systems, or general control systems, as in [24, 33, 23, 10, 11]. In these works, systems were defined in terms of set-valued maps associated with reachable sets (or attainable sets). A similar treatment was also adopted in [29] and related work, where the prolongation sets of reachable sets were used to study stability. In [23], the author established the existence of different types of Lyapunov functions (not necessarily continuous) for both stability and weak stability with respect to closed invariant sets, where “weak stability” means the existence of a stable trajectory from every point outside the invariant set. In [10], the author provided Lyapunov characterizations for both local asymptotic stability and weak asymptotic stability. See [11] for an excellent survey of work along these lines.

It is also possible to reformulate stability for systems with time varying parameters in terms of differential inclusions, as explained earlier; see e.g. [1] and [2]. The first of these books employs Lyapunov functions in

sufficiency characterizations of viability properties (not the same as stability with respect to all solutions), while the second one (see Chapter 6, and especially Section 4) shows various converse theorems that result in nondifferentiable Lyapunov functions, connecting their existence with the solution of optimal control problems. In the recent work [20], one can find conclusions analogous to those in this paper but only for the very special case of linear differential inclusions, resulting in homogeneous “quasiquadratic” Lyapunov functions. Finally, let us mention the work [19] on systems with time varying parameters, in which the author established, under the assumption of *exponential* stability, the existence of differentiable Lyapunov functions on compact sets, for the special case of equilibria.

10. Relations to Stability of Prolongations. In [7, 8, 28, 29, 30], the authors considered various notions of stability for systems of the type (1) (with \mathcal{D} not necessarily compact). These properties are defined in terms of the “prolongations” of the original system. The above papers investigated the relationships between such stability notions and the existence of continuous, not necessarily smooth, Lyapunov functions. In this section, we briefly discuss relations between UGAS stability and the notions considered in those papers, with the purpose of clarifying relations to this related previous work. For the more details on the definitions and elementary properties of prolongation maps and the corresponding stability concepts, we refer the reader to the papers mentioned above.

We start with some abstract definitions. Let $F : \mathbb{R}^n \times \mathbb{R}_{\geq 0} \rightarrow 2^{\mathbb{R}^n}$, $(\xi, t) \mapsto F(\xi, t) \subseteq \mathbb{R}^n$ be any map from $\mathbb{R}^n \times \mathbb{R}_{\geq 0}$ to the set of subsets of \mathbb{R}^n . Associated to F , one defines $\mathfrak{D}F$ and $\mathfrak{J}F$ by

$$\begin{aligned} \mathfrak{D}F(\xi, t) &= \left\{ \eta \in \mathbb{R}^n : \text{there exist sequences } \xi_n, \eta_n \in \mathbb{R}^n, \text{ and } t_n \geq 0 \right. \\ &\quad \left. \text{with } \xi_n \rightarrow \xi, \eta_n \rightarrow \eta, t_n \rightarrow t, \eta \in F(\xi_n, t_n) \right\}, \\ \mathfrak{J}F(\xi, t) &= \left\{ \eta \in \mathbb{R}^n : \text{there exist } t_1, t_2, \dots, t_k \geq 0 \text{ with} \right. \\ &\quad \left. \sum_{i=1}^k t_i = t, \text{ such that } \eta \in F\left(F(\dots F(F(\xi_n, t_1), t_2) \dots, t_{k-1}), t_k\right) \right\}, \end{aligned}$$

where $F(S, t) \stackrel{\text{def}}{=} \bigcup_{\xi \in S} F(\xi, t)$ for any subset S of \mathbb{R}^n .

The map F is called *cluster* if $\mathfrak{D}F = F$, and F is called *transitive* if $\mathfrak{J}F = F$.

For any system (1), consider the reachable set $\mathcal{R}^t(\xi)$ defined in section 5, seen now as a set-valued map. The prolongation map Γ associated with (1) is then defined by letting $\Gamma(\xi, t)$ be the smallest set containing $\mathcal{R}^t(\xi)$ such that Γ is both transitive and cluster. For further discussion regarding the definition of the map Γ , we refer the reader to [28] and the other papers mentioned above.

For subsets A and B of \mathbb{R}^n , we denote the usual distance between the two sets by $d(A, B) = \inf \{d(\xi, \eta) : \xi \in A, \eta \in B\}$. We say that a system (1) is T-stable (we use here the “T” for the name of the author of [28] who, in turn, was inspired by previous work [8]) with respect to a closed, invariant set \mathcal{A} if the following two properties hold:

- There exists a \mathcal{K}_∞ -function $\delta(\cdot)$ such that for any $\varepsilon > 0$,

$$d(\Gamma(\xi, t), \mathcal{A}) < \varepsilon, \quad \text{whenever } |\xi|_{\mathcal{A}} \leq \delta(\varepsilon), \text{ and } t \geq 0;$$

- For any $r, \varepsilon > 0$, there is a $T > 0$ such that

$$d(\Gamma(\xi, t), \mathcal{A}) < \varepsilon, \quad \text{whenever } |\xi|_{\mathcal{A}} < r, \text{ and } t \geq T.$$

Note that this is the same as what is called “global absolute asymptotic stability” (global A.A.S) in [28] for the special case when \mathcal{A} is compact. Clearly, if a system is T-stable, then it is UGAS. It was shown in [28], under some extra technical assumptions, but without the compactness of \mathcal{D} , that global A.A.S implies the existence

of a continuous, not necessarily smooth, Lyapunov function (meaning that V is globally merely continuous; the condition $L_{f_d}V(\xi) \leq -\alpha_3(|\xi|_{\mathcal{A}})$ is replaced by a condition that V should decrease along trajectories).

We will show next that, at least when \mathcal{D} is compact, UGAS implies (and is therefore equivalent to) T-stability. So in what follows in this section, we assume that \mathcal{D} is compact, and also that all systems involved are forward complete. We first need the following fact.

LEMMA 10.1. *For system (1), $\Gamma(\xi, t) = \overline{\mathcal{R}^t(\xi)}$ for any $\xi \in \mathbb{R}^n$ and any $t \geq 0$.*

Proof. First note that the cluster property of Γ implies that $\Gamma(\xi, t)$ is closed for each $\xi \in \mathbb{R}^n$ and each $t \geq 0$. Thus it is enough to show that the map $\mathfrak{R} : (\xi, t) \mapsto \overline{\mathcal{R}^t(\xi)}$ is cluster and transitive.

Take $\xi_0 \in \mathbb{R}^n$ and $\tau > 0$. (The case when $t = 0$ is trivial.) Pick $\eta_0 \in \mathfrak{D}\mathfrak{R}(\xi_0, \tau)$. Then, by definition, there exist sequences $\{\xi_n\}$, $\{\eta_n\}$ and $\{t_n\}$ with $t_n \geq 0$ such that $\xi_n \rightarrow \xi_0$, $\eta_n \rightarrow \eta_0$, $t_n \rightarrow \tau$ and $\eta_n \in \overline{\mathcal{R}^{t_n}(\xi_n)}$.

Note then that for each n , there exists d_n such that

$$|\eta_n - x(t_n, \xi_n, d_n)| < \frac{1}{n}.$$

Let $\zeta_n = x(t_n, \xi_n, d_n)$. Then $\zeta_n \in \mathcal{R}^{t_n}(\xi_n)$ and $\zeta_n \rightarrow \eta_0$. Let K_0 be a compact set such that $\xi_n \in K_0$ for each n , and let $T > 0$ be such that $t_n \leq T$ for any n . Then by Proposition 5.1, there exists a compact set K_1 such that $\mathcal{R}(K_0, T) \subseteq K_1$. Let L be a Lipschitz constant for f with respect to states in K_1 . Then it follows from Gronwall's Lemma that, for n large enough so that $|\xi_n - \xi_0| < e^{-LT}$, it holds that

$$|x(t, \xi_0, d_n) - x(t, \xi_n, d_n)| \leq |\xi_0 - \xi_n|e^{LT},$$

for any $0 \leq t \leq T$. Let $\kappa_n = x(\tau, \xi_0, d_n)$. Then

$$\begin{aligned} |\kappa_n - \zeta_n| &= |x(\tau, \xi_0, d_n) - x(t_n, \xi_n, d_n)| \\ &\leq |x(\tau, \xi_0, d_n) - x(\tau, \xi_n, d_n)| + |x(\tau, \xi_n, d_n) - x(t_n, \xi_n, d_n)| \\ &\leq |\xi_0 - \xi_n|e^{\tau L} + M|\tau - t_n| \end{aligned}$$

where $M = \max\{|f(\xi, \mathbf{d})|, d(\xi, K_1) \leq 1, \mathbf{d} \in \mathcal{D}\}$. It then follows that $\kappa_n \in \mathcal{R}^\tau(\xi_0)$ for each n and $\kappa_n \rightarrow \eta_0$. Thus, we conclude that $\eta_0 \in \overline{\mathcal{R}^\tau(\xi_0)}$. Hence we showed that $\mathfrak{D}\overline{\mathcal{R}^\tau(\xi_0)} = \overline{\mathcal{R}^\tau(\xi_0)}$ for any $\tau > 0$ and any $\xi_0 \in \mathbb{R}^n$, that is, the map \mathfrak{R} is cluster.

To show the transitivity of \mathfrak{R} , first note that, by induction, it is enough to show that

$$(53) \quad \mathfrak{R}(\mathfrak{R}(\xi, t_1), t_2) \subseteq \mathfrak{R}(\xi, t_1 + t_2)$$

for any $\xi \in \mathbb{R}^n$ and any $t_1, t_2 \geq 0$.

Applying Lemma 5.3 to $S = \mathcal{R}^{t_1}(\xi)$ together with the fact that

$$\mathcal{R}^{t_2}(\mathcal{R}^{t_1}(\xi)) = \mathcal{R}^{t_1+t_2}(\xi),$$

one immediately gets (53). \square

Rewriting the definition of UGAS in terms of reachable sets, one has that a system (1) is UGAS if and only if the following properties hold:

- There exists a \mathcal{K}_∞ -function $\delta(\cdot)$ such that for any $\varepsilon > 0$,

$$d(\mathcal{R}^t(\xi), \mathcal{A}) < \varepsilon, \quad \text{whenever } |\xi|_{\mathcal{A}} \leq \delta(\varepsilon), \text{ and } t \geq 0;$$

- For any $r, \varepsilon > 0$, there is a $T > 0$ such that

$$d(\mathcal{R}^t(\xi), \mathcal{A}) < \varepsilon, \quad \text{whenever } |\xi|_{\mathcal{A}} < r, \text{ and } t \geq T.$$

The following conclusion then follows immediately from the continuity of the function $\xi \mapsto d(\xi, \mathcal{A})$ and Lemma 10.1:

PROPOSITION 10.2. *For compact \mathcal{D} , a system (1) is UGAS with respect to \mathcal{A} if and only if it is T-stable. ■*

REMARK 10.3. In the special case when \mathcal{A} is compact, a UGAS system is always forward complete. Thus in that case Proposition 10.2 is still true without completeness. □

REMARK 10.4. The compactness condition on \mathcal{D} is essential. Without the compactness of \mathcal{D} , Proposition 10.2 is in general not true. For instance, the system defined by (50) in section 8 is UGAS with respect to the origin $(0, 0)$. However the system is not T-stable, since $\Gamma(0, t) = \mathbb{R}^2$ for any $t > 0$. Note that for this example, $\overline{R^t(0, t)} = \{0\}$ for any $t > 0$ which is different from $\Gamma(0, t)$. The inconsistency with the conclusion of Lemma 10.1 is caused by the noncompactness of \mathcal{D} . □

Acknowledgements. We wish to thank the Institute for Mathematics and Its Applications for providing an excellent research environment during the Special Year in Control Theory (1992-1993); part of this work was completed while the authors visited the IMA. We also wish to thank John Tsiniás and Randy Freeman for useful comments, and most especially Héctor Sussmann for help with the proof of Proposition 5.1.

REFERENCES

- [1] J.-P. AUBIN, *Viability Theory*, Birkhäuser, Boston, 1991.
- [2] J.-P. AUBIN AND A. CELLINA, *Differential Inclusions: Set-Valued Maps and Viability Theory*, Springer-Verlag, New York, 1984.
- [3] N. P. BHATIA AND G. P. SZEGÖ, *Stability Theory of Dynamical Systems*, Springer-Verlag, New York, 1970.
- [4] T. BRÖCKER, *Differentiable Germs and Catastrophes*, Cambridge University Press, Cambridge, 1975.
- [5] H. FEDERER, *Geometric Measure Theory*, Springer-Verlag, New York, 1969.
- [6] M. W. HIRSCH, *Differential Topology*, Springer-Verlag, New York, 1976.
- [7] N. KALOUPSIDIS, *Accessibility and stability theory of nonlinear control systems*, PhD thesis, Washington University, Saint Louis, Missouri, 1977.
- [8] N. KALOUPSIDIS AND D. L. ELLIOTT, *Stability analysis of the orbits of control systems*, *Mathematical Systems Theory*, 15 (1982), pp. 323–342.
- [9] I. KANELLAKOPOULOS, P. V. KOKOTOVIĆ, AND A. S. MORSE, *A toolkit for nonlinear feedback design*, *Systems & Control Letters*, 18 (1992), pp. 83–92.
- [10] P. E. KLOEDEN, *Eventual stability in general control systems*, *Journal of Differential Equations*, 19 (1975), pp. 106–124.
- [11] ———, *General control systems*, in *Mathematical Control Theory (Proceedings, Canberra, Australia 1977)*, A. Bold and B. Eckmann, eds., Springer-Verlag, New York, 1978, pp. 119–137.
- [12] N. N. KRASOVSKIĬ, *Stability of Motion*, Stanford University Press, Stanford, Ca., 1963.
- [13] J. KURZWEIL, *On the inversion of Lyapunov’s second theorem on stability of motion*, *American Mathematical Society Translations, Series 2*, 24 (1956), pp. 19–77.
- [14] V. LAKSHMIKANTHAM, S. LEELA, AND A. A. MARTYNYUK, *Stability Analysis of Nonlinear Systems*, Marcel Dekker, New York, 1989.
- [15] S. LEFSCHETZ, *Differential Equations: Geometric Theory*, Interscience Publishers, New York, second ed., 1963.
- [16] Y. LIN, *Lyapunov Function Techniques for Stabilization*, PhD thesis, Rutgers, The State University of New Jersey, New Brunswick, New Jersey, 1992.
- [17] Y. LIN, E. D. SONTAG, AND Y. WANG, *Input to state stabilizability for parameterized families of systems*, *International Journal of Robust and Nonlinear Control*, (to appear).
- [18] J. L. MASSERA, *Contributions to stability theory*, *Annals of Mathematics*, 64 (1956), pp. 182–206.
- [19] A. M. MEILAKHS, *Design of stable control systems subject to parametric perturbation*, *Avtomatika i Telemekhanika*, 10 (1978), pp. 5–15.
- [20] A. P. MOLCHANOV AND Y. S. PYANITSKIY, *Criterial of asymptotic stability of differential and difference inclusions encountered in control theory*, *Systems & Control Letters*, 13 (1989), pp. 59–64.
- [21] T. NADZIEJA, *Construction of a smooth Lyapunov function for an asymptotically stable set*, *Czechoslovak Mathematical Journal*, 115 (1990), pp. 195–199.
- [22] L. ROSIER, *Inverse of Lyapunov’s second theorem for measurable functions*, in *Proceedings of Nonlinear Control Systems Design Symposium, Bordeaux, France, June 1992*, M. Fliess, ed., IFAC Publications, 1992, pp. 655–660.

- [23] E. O. ROXIN, *Stability in general control systems*, Journal of Differential Equations, 1 (1965), pp. 115–150.
- [24] P. SEIBERT, *Stability under perturbations in generalized dynamical systems*, in International Symposium on Nonlinear Differential Equations and Nonlinear Mechanics, J. P. LaSalle and S. Lefschetz, eds., New York, 1963, Academic Press, pp. 463–473.
- [25] E. D. SONTAG, *Smooth stabilization implies coprime factorization*, IEEE Transactions on Automatic Control, AC-34 (1989), pp. 435–443.
- [26] ———, *Mathematical Control Theory: Deterministic Finite Dimensional Systems*, Springer-Verlag, New York, 1990.
- [27] E. D. SONTAG AND Y. WANG, *On characterizations of the input-to-state stability property*, Systems & Control Letters, 24 (1995), pp. 351–359. (Preliminary version in “Notions equivalent to input-to-state stability,” in *Proc. IEEE Conf. Decision and Control, Orlando, Dec. 1994*, IEEE Publications, 1994, pp. 3438–3443).
- [28] J. TSINIAS, *A Lyapunov description of stability in control systems*, Nonlinear Analysis, Theory, Methods & Applications, 13 (1989), pp. 63–74.
- [29] J. TSINIAS AND N. KALOUPSIDIS, *Prolongations and stability analysis via lyapunov functions of dynamical polysystems*, Mathematical Systems Theory, 20 (1987), pp. 215–233.
- [30] J. TSINIAS, N. KALOUPSIDIS, AND A. BACCIOTTI, *Lyapunov functions and stability of dynamical polysystems*, Mathematical Systems Theory, 19 (1987), pp. 333–354.
- [31] F. W. WILSON, JR., *Smoothing derivatives of functions and applications*, Transactions of the American Mathematical Society, 139 (1969), pp. 413–428.
- [32] T. YOSHIKAWA, *Stability Theory and the Existence of Periodic Solutions and Almost Periodic Solutions*, Springer-Verlag, New York, 1975.
- [33] V. I. ZUBOV, *Methods of A. M. Lyapunov and Their Application*, P. Noordhoff Ltd., Groningen, The Netherlands, 1964. English edition.

Appendix

A. Some Basic Definitions. In this section we recall some standard concepts from stability theory.

A function $\gamma : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is:

- a \mathcal{K} -function if it is continuous, strictly increasing and $\gamma(0) = 0$;
- a \mathcal{K}_∞ -function if it is a \mathcal{K} -function and also $\gamma(s) \rightarrow \infty$ as $s \rightarrow \infty$;
- a *positive definite* function if $\gamma(s) > 0$ for all $s > 0$, and $\gamma(0) = 0$.

A function $\beta : \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is a \mathcal{KL} -function if:

- for each fixed $t \geq 0$ the function $\beta(\cdot, t)$ is a \mathcal{K} -function, and
- for each fixed $s \geq 0$ it is decreasing to zero as $t \rightarrow \infty$.

Note that we are not requiring β to be continuous in both variables simultaneously; however it turns out in our results that this stronger property will usually hold.

B. Smooth Approximations of Locally Lipschitz Functions. In the proof of the converse Lyapunov theorem, we used a parameterized version of an approximation theorem given in [31]. For convenience of reference, and to make this work self-contained and expository, we next provide the needed variation of the theorem and its proof. (Several details, missing in the proof in [31], have been included as well.)

THEOREM B.1. *Let \mathcal{O} be an open subset of \mathbb{R}^n , and let \mathcal{D} be a compact subset of \mathbb{R}^l , and assume given:*

- a locally Lipschitz function $\Phi : \mathcal{O} \rightarrow \mathbb{R}$;
- a continuous map $f : \mathbb{R}^n \times \mathcal{D} \rightarrow \mathbb{R}^n$, $(x, \mathbf{d}) \mapsto f(x, \mathbf{d})$ which is locally Lipschitz on x uniformly on \mathbf{d} ;
- a continuous function $\alpha : \mathcal{O} \rightarrow \mathbb{R}$ and continuous functions $\mu, \nu : \mathcal{O} \rightarrow \mathbb{R}_{>0}$

such that for each $\mathbf{d} \in \mathcal{D}$,

$$(54) \quad L_{f_{\mathbf{d}}}\Phi(\xi) \leq \alpha(\xi), \quad \text{a.e. } \xi \in \mathcal{O},$$

where $f_{\mathbf{d}}$ is the vector field defined by $f_{\mathbf{d}}(\cdot) = f(\cdot, \mathbf{d})$, (Recall that $\nabla\Phi$ is defined a.e., since Φ is locally Lipschitz by Rademacher’s Theorem, see e.g. [5], page 216). Then there exists a smooth function $\Psi : \mathcal{O} \rightarrow \mathbb{R}$ such that

$$|\Phi(\xi) - \Psi(\xi)| < \mu(\xi), \quad \forall \xi \in \mathcal{O}$$

and for each $\mathbf{d} \in \mathcal{D}$,

$$L_{f_{\mathbf{d}}}\Psi(\xi) \leq \alpha(\xi) + \nu(\xi), \quad \forall \xi \in \mathcal{O}.$$

To prove the theorem, we need first some easy facts about regularization. Let $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth nonnegative function which vanishes outside of the unit disk and satisfies

$$\int_{\mathbb{R}^n} \psi(s) ds = 1.$$

For any measurable, locally essentially bounded function $\Phi : \mathcal{O} \rightarrow \mathbb{R}$ and $0 < \sigma \leq 1$, define the function Φ_σ by convolution with $\frac{1}{\sigma^n} \psi\left(\frac{s}{\sigma}\right)$, that is:

$$(55) \quad \Phi_\sigma(\xi) \stackrel{\text{def}}{=} \int_{\mathbb{R}^n} \Phi(\xi + \sigma s) \psi(s) ds.$$

We think of this function as defined only for those ξ so that $\xi + \sigma s \in \mathcal{O}$ for all $|s| \leq 1$. Note that the integral is finite, as the integrand is essentially bounded and of compact support. The following observation is a standard approximation exercise, so we omit its proof.

LEMMA B.2. *For each compact subset K of \mathcal{O} , there exists some $\sigma_0 > 0$ such that Φ_σ is defined on K , and smooth there, for all $\sigma < \sigma_0$. Moreover, if Φ is continuous, then Φ_σ approaches Φ uniformly on K , as σ tends to 0.* \square

Now assume that Φ is a locally Lipschitz function. Then, for each $\mathbf{d} \in \mathcal{D}$, $L_{f_{\mathbf{d}}}\Phi$ is defined almost everywhere, and furthermore, on any compact subset $K \subseteq \mathcal{O}$,

$$|L_{f_{\mathbf{d}}}\Phi(\xi)| \leq k |f(\xi, \mathbf{d})|, \quad \text{a.e. } \xi \in K, \quad \forall \mathbf{d} \in \mathcal{D},$$

where k is a Lipschitz constant for Φ on K . Therefore, for each \mathbf{d} , (omitting from now on the \mathbb{R}^n in integrals)

$$(L_{f_{\mathbf{d}}}\Phi)_\sigma(\xi) = \int (L_{f_{\mathbf{d}}}\Phi)(\xi + \sigma s) \psi(s) ds$$

is well defined as long as $\xi + \sigma s \in \mathcal{O}$ for all $|s| \leq 1$. Applying Lemma B.2 to $(L_{f_{\mathbf{d}}}\Phi)_\sigma$, this is smooth for any $\sigma > 0$ small.

Suppose that for all $\mathbf{d} \in \mathcal{D}$,

$$(56) \quad L_{f_{\mathbf{d}}}\Phi(\xi) \leq \alpha(\xi), \quad \text{a.e. } \xi \in \mathcal{O},$$

for some continuous function α . Pick any compact subset $K \subseteq \mathcal{O}$. On this set K , we have

$$\begin{aligned} (L_{f_{\mathbf{d}}}\Phi)_\sigma(\xi) &= \int (L_{f_{\mathbf{d}}}\Phi)(\xi + \sigma s) \psi(s) ds \leq \int \alpha(\xi + \sigma s) \psi(s) ds \\ &\leq \alpha(\xi) + \max_{|s| \leq 1, \xi \in K} |\alpha(\xi + \sigma s) - \alpha(\xi)|. \end{aligned}$$

From here we get the following conclusion:

LEMMA B.3. *For any compact subset K of \mathcal{O} , $(L_{f_{\mathbf{d}}}\Phi)_\sigma$ is a C^∞ function defined on K for all σ small enough, and, if (56) holds for all $\mathbf{d} \in \mathcal{D}$ and all $\xi \in \mathcal{O}$, then for any $\varepsilon > 0$ given, there exists some $\sigma_0 > 0$ such that*

$$(L_{f_{\mathbf{d}}}\Phi)_\sigma(\xi) \leq \alpha(\xi) + \varepsilon$$

for all $\sigma \leq \sigma_0$, all $\mathbf{d} \in \mathcal{D}$, and all $\xi \in K$. \square

The following lemma illustrates the relationship between $L_{f_{\mathbf{d}}}(\Phi_\sigma)$ and $(L_{f_{\mathbf{d}}}\Phi)_\sigma$.

LEMMA B.4. *On any compact subset K of \mathcal{O} ,*

$$\sup_{\mathbf{d} \in \mathcal{D}, \xi \in K} |L_{f_{\mathbf{d}}}(\Phi_{\sigma})(\xi) - (L_{f_{\mathbf{d}}}\Phi)_{\sigma}(\xi)| \rightarrow 0$$

as σ tends to 0.

Proof. For each $\xi \in \mathcal{O}$, we use $\varphi(t, \xi, \mathbf{d})$ to denote the solution of the differential equation:

$$\dot{x} = f(x, \mathbf{d})$$

with the initial condition $\varphi(0, \xi, \mathbf{d}) = \xi$. It follows from the assumptions on f and compactness of K and \mathcal{D} that there exist some compact neighborhood V of K and some $\tau_1 > 0$ and $\sigma_0 > 0$ such that $\varphi(t, \xi + \sigma s, \mathbf{d}) \in V$ for all $\xi \in K$, $|s| \leq 1$, $\sigma \leq \sigma_0$, $\mathbf{d} \in \mathcal{D}$ and $|t| \leq \tau_1$.

For the Lipschitz function Φ , we have, for all ξ, \mathbf{d} and $\sigma \leq \sigma_0$:

$$\begin{aligned} L_{f_{\mathbf{d}}}(\Phi_{\sigma})(\xi) &= \left. \frac{d}{dt} \right|_{t=0} \Phi_{\sigma}(\varphi(t, \xi, \mathbf{d})) = \left. \frac{d}{dt} \right|_{t=0} \int \Phi(\varphi(t, \xi, \mathbf{d}) + \sigma s) \psi(s) ds \\ &= \lim_{t \rightarrow 0} \frac{1}{t} \int (\Phi(\varphi(t, \xi, \mathbf{d}) + \sigma s) - \Phi(\xi + \sigma s)) \psi(s) ds, \end{aligned}$$

and

$$(57) \quad (L_{f_{\mathbf{d}}}\Phi)_{\sigma}(\xi) = \int L_{f_{\mathbf{d}}}\Phi(\xi + \sigma s) \psi(s) ds$$

$$(58) \quad = \int \left. \frac{d}{dt} \right|_{t=0} \Phi(\varphi(t, \xi + \sigma s, \mathbf{d})) \psi(s) ds$$

$$(59) \quad = \lim_{t \rightarrow 0} \frac{1}{t} \int [\Phi(\varphi(t, \xi + \sigma s, \mathbf{d})) - \Phi(\xi + \sigma s)] \psi(s) ds.$$

Notice that the integrand in (57) equals that in (58) almost everywhere on s (for each fixed ξ and σ) and that (59) follows from (58) because of the Lebesgue Dominated Convergence Theorem and the following fact:

$$\begin{aligned} &\frac{1}{|t|} |\Phi(\varphi(t, \xi + \sigma s, \mathbf{d})) - \Phi(\xi + \sigma s)| \psi(s) \\ &\leq \frac{k}{|t|} |\varphi(t, \xi + \sigma s, \mathbf{d}) - (\xi + \sigma s)| \psi(s) \leq kC\psi(s), \quad \forall t \in [-\tau_1, \tau_1], \end{aligned}$$

where $C \stackrel{\text{def}}{=} \max_{\xi \in V, \mathbf{d} \in \mathcal{D}} |f(\xi, \mathbf{d})|$ and k is a Lipschitz constant for Φ on V .

Now one sees that

$$L_{f_{\mathbf{d}}}(\Phi_{\sigma})(\xi) - (L_{f_{\mathbf{d}}}\Phi)_{\sigma}(\xi) = \lim_{t \rightarrow 0} \frac{1}{t} \int [\Phi(\varphi(t, \xi, \mathbf{d}) + \sigma s) - \Phi(\varphi(t, \xi + \sigma s, \mathbf{d}))] \psi(s) ds.$$

Thus it is enough to show that for any $\varepsilon > 0$, there exist some $\delta > 0$ and $\tau^* > 0$ such that the above integral is bounded by ε for all $\mathbf{d} \in \mathcal{D}$, $\xi \in K$, $|t| < \tau^*$ and $\sigma < \delta$. This is basically a standard argument on continuous dependence on initial conditions, but we provide the details. For $0 \leq \tau \leq \tau_1$, let

$$\gamma(\tau) \stackrel{\text{def}}{=} \sup \{ |f(\varphi(t, \zeta, \mathbf{d}), \mathbf{d}) - f(\zeta, \mathbf{d})| : |t| \leq \tau, \zeta \in V, \mathbf{d} \in \mathcal{D} \}.$$

Then $\gamma(0) = 0$, and γ is nondecreasing and continuous at $t = 0$, because

$$|f(\varphi(t, \zeta, \mathbf{d}), \mathbf{d}) - f(\zeta, \mathbf{d})| \leq C_3 |\varphi(t, \zeta, \mathbf{d}) - \zeta| \leq C_3 C_4 |t|,$$

where C_3 is a (uniform) Lipschitz constant for f on V_1 , C_4 is an upper bound for $|f(\xi, \mathbf{d})|$ on V_1 , and V_1 is some compact neighborhood of V such that $\varphi(t, \zeta, \mathbf{d}) \in V_1$ for any $\zeta \in V$, $\mathbf{d} \in \mathcal{D}$ and $|t| \leq \tau_1$. For any $\zeta \in V$, $\mathbf{d} \in \mathcal{D}$ and $|t| \leq \tau_1$,

$$|\varphi(t, \zeta, \mathbf{d}) - (\zeta + tf(\zeta, \mathbf{d}))| \leq \int_0^{|t|} \gamma(\tau) d\tau \leq |t| \gamma(|t|).$$

Now for $\xi \in K$, we have

$$\begin{aligned} & |\Phi(\varphi(t, \xi, \mathbf{d}) + \sigma s) - \Phi(\varphi(t, \xi + \sigma s, \mathbf{d}))| \leq k |\varphi(t, \xi, \mathbf{d}) + \sigma s - \varphi(t, \xi + \sigma s, \mathbf{d})| \\ & \leq k |\xi + \sigma s + tf(\xi, \mathbf{d}) - (\xi + \sigma s + tf(\xi + \sigma s, \mathbf{d}))| \\ & \quad + k |\varphi(t, \xi, \mathbf{d}) - (\xi + tf(\xi, \mathbf{d}))| + k |\varphi(t, \xi + \sigma s, \mathbf{d}) - (\xi + \sigma s + tf(\xi + \sigma s, \mathbf{d}))| \\ (60) \quad & \leq k |t| |f(\xi, \mathbf{d}) - f(\xi + \sigma s, \mathbf{d})| + 2k |t| \gamma(|t|). \end{aligned}$$

Finally, for $\varepsilon > 0$, let δ and τ^* be such that

$$\gamma(\tau) < \frac{\varepsilon}{3k} \quad \text{and} \quad |f(\xi, \mathbf{d}) - f(\xi + \sigma s, \mathbf{d})| < \frac{\varepsilon}{3k}$$

for any $\xi \in K$, $\mathbf{d} \in \mathcal{D}$, $|s| \leq 1$, $\sigma < \delta$ and $|t| < \tau^*$. It then follows from (60) that

$$\frac{1}{|t|} \int [\Phi(\varphi(t, \xi, \mathbf{d}) + \sigma s) - \Phi(\varphi(t, \xi + \sigma s, \mathbf{d}))] \psi(s) ds < \int \varepsilon \psi(s) ds = \varepsilon$$

for any $\xi \in K$, $\mathbf{d} \in \mathcal{D}$, $|t| < \tau^*$ and $\sigma < \delta$, which implies

$$|L_{f_{\mathbf{d}}}(\Phi_{\sigma})(\xi) - (L_{f_{\mathbf{d}}}\Phi)_{\sigma}(\xi)| < \varepsilon$$

for any $\sigma < \sigma_0$, $\mathbf{d} \in \mathcal{D}$ and $\xi \in K$. \square

Combining the previous three lemmas, we obtain the following conclusion:

LEMMA B.5. *Let K be a compact subset of \mathcal{O} . Then for any given $\varepsilon > 0$, there exists some smooth function Ψ defined on K such that*

$$|\Psi(\xi) - \Phi(\xi)| < \varepsilon \quad \text{and} \quad L_{f_{\mathbf{d}}}\Psi(\xi) \leq \alpha(\xi) + \varepsilon$$

for all $\xi \in K$, $\mathbf{d} \in \mathcal{D}$. \square

Now we are ready to complete the proof of Theorem (B.1). For the open subset \mathcal{O} of \mathbb{R}^n , let $\{\mathcal{U}_i\}$ be a locally finite countable cover of \mathcal{O} with $\bar{\mathcal{U}}_i$ compact and $\bar{\mathcal{U}}_i \subseteq \mathcal{O}$. Let $\{\beta_i\}$ be a partition of unity on \mathcal{O} subordinate to $\{\mathcal{U}_i\}$. For any given positive functions $\mu(\cdot)$ and $\nu(\cdot)$, let

$$\varepsilon_i \stackrel{\text{def}}{=} \min \left\{ \inf_{\xi \in \mathcal{U}_i} \mu(\xi), \inf_{\xi \in \mathcal{U}_i} \nu(\xi) \right\}.$$

For each i , it follows from Lemma B.5 that there exists some smooth function Ψ_i defined on $\bar{\mathcal{U}}_i$ such that

$$|\Phi(\xi) - \Psi_i(\xi)| < \frac{\varepsilon_i}{2^{i+1}(1 + \tau_i)} \quad \text{and} \quad L_{f_{\mathbf{d}}}\Psi_i(\xi) \leq \alpha(\xi) + \frac{\varepsilon_i}{2}$$

on $\bar{\mathcal{U}}_i$, where $\tau_i \stackrel{\text{def}}{=} \max\{|L_{f_{\mathbf{d}}}\beta_i(\xi)| : \xi \in \bar{\mathcal{U}}_i, \mathbf{d} \in \mathcal{D}\}$. We define $\Psi = \sum_i \beta_i \Psi_i$. Clearly Ψ is a smooth function defined on \mathcal{O} , and

$$\begin{aligned} |\Psi(\xi) - \Phi(\xi)| & \leq \sum_{j \in \mathcal{J}_{\xi}} \beta_j(\xi) |\Psi_j(\xi) - \Phi(\xi)| \\ & < \max_{j \in \mathcal{J}_{\xi}} \varepsilon_j \leq \mu(\xi), \end{aligned}$$

where $\mathcal{J}_\xi \stackrel{\text{def}}{=} \{j : \xi \in \mathcal{U}_j\}$.
 For $L_{f_d}\Psi$, one has

$$\begin{aligned}
 L_{f_d}\Psi(\xi) &= L_{f_d}\Phi(\xi) + L_{f_d}\left(\sum_i \beta_i(\Psi_i - \Phi)\right)(\xi) \\
 &= L_{f_d}\Phi(\xi) + \sum (L_{f_d}\beta_i)(\Psi_i - \Phi)(\xi) + \sum \beta_i(L_{f_d}\Psi_i(\xi) - L_{f_d}\Phi(\xi)) \\
 &= \sum_{j \in \mathcal{J}_\xi} (L_{f_d}\beta_j)(\Psi_j - \Phi)(\xi) + \sum_{j \in \mathcal{J}_\xi} \beta_j L_{f_d}\Psi_j(\xi) \\
 &< \sum_{j \in \mathcal{J}_\xi} \frac{\varepsilon_j}{2^{j+1}} + \sum_{j \in \mathcal{J}_\xi} \beta_j(\xi) \left(\alpha(\xi) + \frac{\varepsilon_i}{2} \right) \\
 &\leq \frac{1}{2} \max_{j \in \mathcal{J}_\xi} \{\varepsilon_j\} + \alpha(\xi) + \frac{1}{2} \max_{j \in \mathcal{J}_\xi} \{\varepsilon_j\} \\
 &\leq \alpha(\xi) + \nu(\xi).
 \end{aligned}$$

We conclude that Ψ is the desired function. ■