

**BOTT PERIODICITY FOR GROUP RINGS
AN APPENDIX TO
“PERIODICITY OF HERMITIAN K -GROUPS”**

CHARLES WEIBEL

ABSTRACT. We show that the groups $K_n(RG; \mathbb{Z}/m)$ are Bott-periodic for $n \geq 1$ whenever G is a finite group, m is prime to $|G|$, R is a ring of S -integers in a number field and $1/m \in R$.

For any positive integer m there is a Bott element $b_K \in K_p(\mathbb{Z}[1/m]; \mathbb{Z}/m)$, where the period $p = p(m)$ is: $2(\ell - 1)\ell^{\nu-1}$ if $m = \ell^\nu$ and ℓ is an odd prime; $\max\{8, 2^{\nu-1}\}$ if $m = 2^\nu$; and $\prod p(m_i)$ if $m = \prod m_i$ is the factorization of m into primary factors.

In this appendix we consider a finite group G of order prime to m , and consider the Bott periodicity map $x \mapsto x \cdot b_K$ from $K_n(R[G]; \mathbb{Z}/m)$ to $K_{n+p}(R[G]; \mathbb{Z}/m)$ for rings of integers R in local and global fields.

Theorem 0.1. *Assume that m is relatively prime to $|G|$, and that R is a ring of S -integers in a number field F with $1/m \in R$. Then the Bott periodicity maps $b_K : K_n(R[G]; \mathbb{Z}/m) \rightarrow K_{n+p}(R[G]; \mathbb{Z}/m)$ are isomorphisms for all $n \geq 1$.*

Theorem 0.2. *Assume that m is relatively prime to $|G|$, and that R is the ring of integers in a local field F with $1/m \in R$. Then the Bott periodicity maps $b_K : K_n(R[G]; \mathbb{Z}/m) \rightarrow K_{n+p}(R[G]; \mathbb{Z}/m)$ are isomorphisms for all $n \geq 0$,*

Theorems 0.1 and 0.2 are used in [BKO] to show that $KQ_n(R[G]; \mathbb{Z}/m)$ also satisfies Bott periodicity. Since this is immediate if m is odd, when the non-Witt part of $KQ_n(A; \mathbb{Z}/m)$ is a summand of $K_n(A; \mathbb{Z}/m)$, this result is primarily interesting for m even and G a (solvable) group of odd order.

Remark. The proofs show that we may replace $R[G]$ by any order in $F[G]$.

The oldest result of this kind is due to Browder, who proved in [1, 2.6] that the Bott periodicity map b_K is an isomorphism for finite fields and $n \geq 0$ (when m is prime, which implies periodicity for all m).

Almost as old is the following folklore result, which includes finite group rings.

Lemma 0.3. *If B is a finite ring and $1/m \in B$, then the Bott periodicity map $K_n(B, \mathbb{Z}/m) \rightarrow K_{n+p}(B, \mathbb{Z}/m)$ is an isomorphism for all $n \geq 0$.*

Proof. If \mathfrak{m} is the nilradical of B , then $B_{\text{red}} = B/\mathfrak{m}$ is semisimple. As such it is a product of matrix rings over finite fields. Now $K_*(B; \mathbb{Z}/m) \cong K_*(B_{\text{red}}; \mathbb{Z}/m)$ by [9, 1.4]. By Morita invariance, we are reduced to the Browder’s theorem that the Bott periodicity map is an isomorphism for finite fields. \square

Remark 0.4. *The finite groups $K_n(\mathbb{F}_2[C_2]; \mathbb{Z}/8)$ were computed by Hesselholt and Madsen in [4]; they are not Bott periodic as their order goes to infinity with n .*

The next step is to consider the semisimple group ring $F[G]$ when F is a number field. We will use the fact that multiplication by b_K is an isomorphism on

$K_n(F; \mathbb{Z}/m)$ for all $n \geq 1$. This is a consequence of the Milnor-Bloch-Kato conjecture (see [6] [12] [11]), and has been observed by several people.

Lemma 0.5. *If F is a number field and m is prime to $|G|$, the Bott periodicity map $b_K : K_n(F[G]; \mathbb{Z}/m) \rightarrow K_{n+p}(F[G]; \mathbb{Z}/m)$ is an isomorphism for all $n \geq 1$.*

Proof. By Maschke's Theorem, $F[G]$ is the product of simple rings $A_i = M_{n_i}(D_i)$. The *Schur index* of the division algebra D_i is the integer r_i such that D_i has dimension r_i^2 over its center F_i . By [2, 27.11], each $n_i r_i$ divides $|G|$ and so is prime to m . Now the composite of the reduced norm $N_{red} : K_*(D_i; \mathbb{Z}/m) \rightarrow K_*(F_i; \mathbb{Z}/m)$ and inclusion $\iota^* : K_*(F_i; \mathbb{Z}/m) \rightarrow K_*(D_i; \mathbb{Z}/m)$ is multiplication by r_i in either direction, so these are isomorphisms. Since the Bott periodicity map commutes with ι^* , and is an isomorphism on $K_n(F_i; \mathbb{Z}/m)$ for $n \geq 1$, it is an isomorphism on each factor $K_n(A_i; \mathbb{Z}/m)$ in this range. \square

Lemma 0.6. *Let Λ be a maximal R -order in a simple factor A of $F[G]$. Then the periodicity map $b_K : K_n(\Lambda; \mathbb{Z}/m) \rightarrow K_{n+p}(\Lambda; \mathbb{Z}/m)$ is an isomorphism for $n \geq 1$.*

Proof. Let R_E be the integral closure of R in the center E of A . For each prime ideal \mathfrak{p} of R_E the ring $\Lambda/\mathfrak{p}\Lambda$ is a finite ring. Since the Bott element commutes with the localization sequence for $\Lambda \rightarrow A$ we have a commutative diagram (with finite coefficients omitted):

$$\begin{array}{ccccccccc} K_{n+1}(A) & \rightarrow & \oplus K_n(\Lambda/\mathfrak{p}\Lambda) & \rightarrow & K_n(\Lambda) & \rightarrow & K_n(A) & \rightarrow & \oplus K_{n-1}(\Lambda/\mathfrak{p}\Lambda) \\ \simeq \downarrow b_K & & \simeq \downarrow b_K & & \downarrow b_K & & \simeq \downarrow b_K & & \simeq \downarrow b_K \\ K_{n+p+1}(A) & \rightarrow & \oplus K_{n+p}(\Lambda/\mathfrak{p}\Lambda) & \rightarrow & K_{n+p}(\Lambda) & \rightarrow & K_{n+p}(A) & \rightarrow & \oplus K_{n+p-1}(\Lambda/\mathfrak{p}\Lambda). \end{array}$$

By Lemmas 0.5 and 0.3, the four outer verticals are isomorphisms for $n \geq 1$. By the 5-lemma, the middle vertical map is an isomorphism for $n \geq 1$. \square

Proof of Theorem 0.1. As noted in [8, 6.5], the groups $NK_*(R[G]; \mathbb{Z}/m)$ are zero because $m \nmid |G|$. Therefore the spectral sequence

$$E_{p,q}^1 = N^p K_q(R[G]; \mathbb{Z}/m) \Rightarrow KH_{p+q}(R[G]; \mathbb{Z}/m)$$

(cf. [10, 1.6]) degenerates to yield

$$K_*(R[G]; \mathbb{Z}/m) \cong KH_*(R[G]; \mathbb{Z}/m).$$

Now $R[G]$ is an order in the semisimple algebra $F[G]$, and is contained in a maximal order Λ . Note that Λ decomposes as a product of maximal orders Λ_i in the simple factors A_i of $F[G]$. Let I be the conductor ideal from the maximal order Λ to $R[G]$; the quotient rings $R[G]/I$ and Λ/I are finite. Since the Bott element $b_K \in K_P(\mathbb{Z}[1/m]; \mathbb{Z}/m)$ is compatible with the Mayer-Vietoris sequences [10, 2.2], we have a commutative diagram, where B denotes $\Lambda \times (R[G]/I)$ and C denotes Λ/I :

$$\begin{array}{ccccccccc} KH_{n+1}(B) & \rightarrow & KH_{n+1}(C) & \rightarrow & KH_n(R[G]) & \rightarrow & KH_n(B) & \rightarrow & KH_n(C) \\ \simeq \downarrow b_K & & \simeq \downarrow b_K & & \downarrow b_K & & \simeq \downarrow b_K & & \simeq \downarrow b_K \\ KH_{n+p+1}(B) & \rightarrow & KH_{n+p+1}(C) & \rightarrow & KH_{n+p}(R[G]) & \rightarrow & KH_{n+p}(B) & \rightarrow & KH_{n+p}(C) \end{array}$$

When $n \geq 1$, the outer periodicity maps b_K are isomorphisms by Lemmas 0.6 and 0.3. Hence the middle map b_K is also an isomorphism for $n \geq 1$. \square

We now consider the case of local fields.

Theorem 0.7. *Let R be the ring of integers in a local field F with $1/m \in R$. If Λ is a maximal order in a finite-dimensional central simple F -algebra A , then the periodicity maps $K_n(\Lambda; \mathbb{Z}/m) \rightarrow K_{n+p}(\Lambda; \mathbb{Z}/m)$, resp., $K_n(A; \mathbb{Z}/m) \rightarrow K_{n+p}(A; \mathbb{Z}/m)$ are isomorphisms for all $n \geq 0$, resp., for all $n \geq 1$.*

Proof. There is a division algebra D so that $A \cong M_r(D)$ and an inner automorphism of A identifying Λ with $M_r(\Delta)$, where Δ is the unique maximal order in D ; see [2, 26.23]. Let B be the finite residue matrix algebra $M_r(\bar{\Delta})$ of A . By Suslin-Yufryakov [7], $K_n(\Lambda; \mathbb{Z}/m) \cong K_n(B; \mathbb{Z}/m)$. By Lemma 0.3, the Bott map is an isomorphism on $K_n(\Lambda; \mathbb{Z}/m)$ for all $n \geq 0$. Since the Bott element acts naturally on the localization sequence $K_*(B; \mathbb{Z}/m) \rightarrow K_*(\Lambda; \mathbb{Z}/m) \rightarrow K_*(A; \mathbb{Z}/m)$, it follows that the $K_n(A; \mathbb{Z}/m)$ are also Bott periodic for $n \geq 1$. \square

Proof of Theorem 0.2. The proof of Theorem 0.1 goes through, replacing Lemma 0.6 with Theorem 0.7. \square

REFERENCES

- [BKO] A. J. Berrick, M. Karoubi and P. A. Østvær, *Periodicity of hermitian K -groups*, preprint, 2009.
- [1] W. Browder, *Algebraic K -theory with coefficients \mathbb{Z}/p* , pp. 40–84 in *Lecture Notes in Math.* 657, Springer-Verlag, 1978.
- [2] C. Curtis and I. Reiner, *Methods of Representation Theory*, Wiley & Sons, New York, 1981.
- [3] O. Gabber, *K -theory of Henselian local rings and Henselian pairs*, *AMS Contemp. Math.* 126 (1992), 59–70.
- [4] L. Hesselholt and I. Madsen, *Cyclic polytopes and the K -theory of truncated polynomial algebras*, *Invent. Math.* 130 (1997), 73–97.
- [5] J. Hornbostel and S. Yagunov, *Rigidity for Henselian local rings and \mathbb{A}^1 -representable theories*, *Math. Zeit.* 255 (2007), 437–449.
- [6] V. Voevodsky, *Motivic cohomology with $\mathbb{Z}/2$ coefficients*, *Publ. Math. Inst. Hautes Études Sci.* 98 (2003), 59–104.
- [7] A. Suslin and A. Yufryakov, *The K -theory of local division algebras*. (Russian) *Dokl. Akad. Nauk SSSR* 288 (1986), 832–836. English Transl. *Soviet Math. Dokl.* 33 (1986), 794–798.
- [8] C. Weibel, *Mayer-Vietoris sequences and module structures on NK_** , pp. 466–493 in *Lecture Notes in Math.*, #854, Springer-Verlag, 1981.
- [9] C. Weibel, *Mayer-Vietoris sequences and mod p K -theory*, pp. 390–407 in *Lecture Notes in Math.*, #966, Springer-Verlag, 1982.
- [10] C. Weibel, *Homotopy algebraic K -theory*, *AMS Contemp Math.*, **83** (1989), 461–488.
- [11] C. Weibel, *Algebraic K -theory of rings of integers in local and global fields*, chapter 5 (pp. 139–184) in *Handbook of K -theory*, Springer-Verlag, 2005. Archived at <http://math.uiuc.edu/K-theory/0691>
- [12] C. Weibel, *The Norm Residue Isomorphism Theorem*, *J. Topology* 2 (2009), 346–372.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854
E-mail address, Charles Weibel: weibel@math.rutgers.edu