

ON THE MONOCHROMATIC SCHUR TRIPLES TYPE PROBLEM

THOTSAPORN “AEK” THANATIPANONDA

ABSTRACT. We discuss a problem posed by Ronald Graham about the minimum number, over all 2-colorings of $[1, n]$, of monochromatic $(x, y, x + ay)$ triples for $a \geq 1$. We give a new proof of the original case of $a = 1$. We show that the minimum number of such triples is at most $\frac{n^2}{2a(a^2+2a+3)} + O(n)$ when $a \geq 2$. We also find a new upper bound for the minimum number, over all r -colorings of $[1, n]$, of monochromatic Schur triples, for $r \geq 3$.

1. INTRODUCTION

The *Schur numbers*, $s(r)$, denote the maximal integer n such that there exists an r -coloring of $[1, n - 1]$ that avoids a monochromatic solution to $x + y = z$. For example $s(2) = 5$ and $s(3) = 14$. $s(5)$ is unknown but is conjectured to be 161.

The original question about the minimum number, over all 2-colorings of $[1, n]$, of monochromatic Schur triples was asked by Ronald Graham in 1997. It can be thought of as a bigger scale version of Schur numbers. It was solved in 1998. The answer is $\frac{n^2}{22} + O(n)$ that is realized by coloring the first $\frac{4n}{11}$ integers red, the next $\frac{6n}{11}$ integers blue, and the final $\frac{n}{11}$ integers red. The first two solutions were given by Robertson and Zeilberger [4] and Schoen [5]. Later Datskovsky [1] found another proof.

Ronald Graham asked another question generalizing the original one. The question was about the minimum number of monochromatic $(x, y, x + ay)$ triples, $a \geq 2$ on $[1, n]$. We discuss this problem in this chapter.

1991 *Mathematics Subject Classification*. 05D10.

Key words and phrases. Ramsey Theory.

In section 2, we give a new simple proof of the original problem of finding the minimum number, over all 2-colorings of $[1, n]$, of monochromatic Schur triples. In section 3, we talk about the generalized problem asked by Graham. For this problem, we wrote a computer program to find an optimal coloring for small n to see some patterns. Then we used a newly found “greedy calculus” to obtain a “good” upper bound. The final step was to try to match the lower bound and upper bound of the problem. In section 4, we also apply the greedy calculus to the original question on Schur triples with $r \geq 3$, to obtain a new upper bound.

2. THE MINIMUM NUMBER, OVER ALL 2-COLORINGS OF $[1, n]$, OF MONOCHROMATIC SCHUR TRIPLES

2.1 A Greedy Algorithm for The Upper bound

It is natural to find examples of good colorings first. This example will give us an upper bound. Then we try to show that this upper bound is also a lower bound.

We will show how to find an upper bound for the minimum number, over all 2-colorings of $[1, n]$, of monochromatic triples that are solutions of $x + y = z$. We will obtain this upper bound by using the Greedy Algorithm. We denote the colors red and blue.

The general idea is to keep adding more new intervals with different colors so that, each time, the overall coloring has the least number of monochromatic triples. For other proofs of this original problem see [4], [1], [5].

First

We paint the first interval of length k red. We will have $\frac{k^2}{4} + O(k)$ monochromatic triples solution of $x + y = z$.

Second

We paint the second interval blue. We want to find the length of the interval (with this color) so that the overall number of the monochromatic triples is minimized.

Let the length of this interval be $(1 + j)k$ (here j is the number we want to find).

The total number of monochromatic triples on the whole interval is now $\frac{k^2}{4} + \frac{j^2k^2}{4} = \frac{(1+j^2)k^2}{4}$.

The total length is $N = k + (1 + j)k = (2 + j)k$.

So the total number of monochromatic triples in terms of N is

$$\frac{(1+j^2)(\frac{N}{2+j})^2}{4} = \frac{(1+j^2)N^2}{(2+j)^2 4}.$$

To find the minimum, we use calculus to get $j = \frac{1}{2}$. The total number of monochromatic Schur triples is then $\frac{N^2}{20} + O(n)$.

So far so good. We have a coloring that paints the first k integers red, followed by painting the next $(1 + \frac{1}{2})k$ integers blue.

Third

Now we try to stick red at the end of the interval, and try to lower the overall number of triples. Say the length of this interval is jk , where j is the number we want to find.

The total number of monochromatic Schur triples on the whole interval is $\frac{k^2}{4} + \frac{k^2}{16} + \frac{j^2k^2}{2} = (\frac{5}{16} + \frac{j^2}{2})k^2$.

The total length is $N = k + (1 + \frac{1}{2})k + jk = (\frac{5}{2} + j)k$.

So the total number of monochromatic Schur triples in term of N is

$$(\frac{5}{16} + \frac{j^2}{2})\frac{N^2}{(\frac{5}{2}+j)^2} = \frac{5+8j^2}{(5+2j)^2} \frac{N^2}{4}.$$

To find the minimum, we again use calculus and get $j = \frac{1}{4}$. The total number of monochromatic triples is $\frac{N^2}{22} + O(n)$. The coloring for the whole interval is a red interval of length equal to k , a blue interval of length equal to $(1 + \frac{1}{2})k$ and another red interval of length equal to $\frac{1}{4}k$. k is such that the sum of these intervals is N , i.e. $k = \frac{N}{(\frac{5}{2}+\frac{1}{4})} = \frac{4N}{11}$.

Fourth

We try to lower the bound further by having a blue interval of length, say, jk at the end of the previous interval. But now we get that the minimizing j is negative. So we stop.

As a conclusion, the optimal coloring is proportional to $[1, \frac{3}{2}, \frac{1}{4}]$, with colors $[R, B, R]$ yielding that indeed the minimal number is $\frac{N^2}{22} + O(n)$.

2.2 The Lower Bound

Finding lower bound is, in general, the difficult part. However, in this case, it is possible since we can turn the problem into a calculus problem. A similar technique was used in [3]

Definition

Let $M_\chi(n)$ be the number of monochromatic Schur triples for a 2-coloring χ of $[1, n]$.

Let Q be two times the number of non-monochromatic Schur triples for a 2-coloring of $[1, n]$.

Divide the interval $[1, n]$ into k consecutive intervals.

Let r_i be the number of red points in the interval I_i .

Let b_i be the number of blue points in the interval I_i .

Let $S_{i,j}$ be the number of non-monochromatic pairs in the square of $I_i \times I_j$.

Let $T_{i,j}$ be the number of non-monochromatic pairs in the triangle of $I_i \times I_j$.

Note: $r_i + b_i = \frac{n}{k}$.

Lemma 1) $M_\chi(n) = \frac{n^2}{4} - \frac{Q}{2} + O(n)$.

The total number of triples = |monochromatic triples| + |non-monochromatic triples|
 $= M_\chi(n) + \frac{Q}{2}$.

Since the total number of triples is $\frac{n^2}{4} + O(n)$, we have $M_\chi(n) = \frac{n^2}{4} - \frac{Q}{2} + O(n)$. \square

The plan is to find an upper bound of Q that will give the lower bound of $M_\chi(n)$.

Lemma 2) $Q = |R||B| + \frac{1}{2}(\sum_{i+j < k} S_{i,j} + \sum_{i=1..k} T_{i,k-i+1})$,
 where $|R| = \sum_{i=1..k} r_i$ and $|B| = \sum_{i=1..k} b_i$.

$$\begin{aligned} Q &= |\{(R, B), (B, R) \mid y - x \geq 0\}| + |\{(R, B), (B, R) \mid x + y \leq n, x \geq y\}| \\ &= |\{(R, B), (B, R) \mid y - x \geq 0\}| + \frac{1}{2} |\{(R, B), (B, R) \mid x + y \leq n\}|. \end{aligned}$$

Note that each non-monochromatic triple contributes two non-monochromatic pairs: for example, $(x, y, z) = (R, B, R)$ gives $(x, y) = (R, B)$ and $(y, z) = (B, R)$. The statement of the lemma follows. \square .

Now we find an upper bound for Q .

For each $T_{i,j}$ we have two ways to bound it:

- 1) $T_{i,j} \leq \text{area of the triangle} = \frac{1}{2} \binom{n}{k}^2$.
- 2) $T_{i,j} \leq S_{i,j}$.

Example 1: $k = 2$, with the upper bound of $T_{1,2}, T_{2,1}$ using the areas of the triangles. We have

$$\begin{aligned} Q &= |R| |B| + \frac{1}{2}(S_{1,1} + T_{1,2} + T_{2,1}) \\ &\leq (r_1 + r_2)(b_1 + b_2) + r_1 b_1 + \frac{n^2}{8} \\ &= (r_1 + r_2)(n - r_1 - r_2) + r_1 \left(\frac{n}{2} - r_1\right) + \frac{n^2}{8}. \end{aligned}$$

We use calculus to find a maximum of Q where $0 \leq r_1, r_2 \leq \frac{n}{2}$. The optimal solutions is $r_1 = \frac{n}{4}$ and $r_2 = \frac{n}{4}$.

We then get the maximum Q as $\frac{7n^2}{16}$. This yields $M_\chi(n) \geq \frac{n^2}{32} + O(n)$. \square

Example 2: $k = 3$, with the upper bound of $T_{1,3}, T_{3,1}$ using the areas of the triangles. The upper bound of $T_{2,2}$ using $S_{2,2}$. We have

$$\begin{aligned} Q &= |R| |B| + \frac{1}{2}(S_{1,1} + S_{1,2} + S_{2,1} + T_{1,3} + T_{2,2} + T_{3,1}) \\ &\leq (r_1 + r_2 + r_3)(b_1 + b_2 + b_3) + r_1 b_1 + r_1 b_2 + r_2 b_1 + r_2 b_2 + \frac{n^2}{18}. \end{aligned}$$

We use calculus to find a maximum of Q where $0 \leq r_1, r_2, r_3 \leq \frac{n}{3}$. One of the optimal solution is $r_1 = 0, r_2 = \frac{n}{3}$ and $r_3 = \frac{n}{6}$.

This yields the maximum Q is $\frac{5n^2}{12}$ which leads to $M_\chi(n) \geq \frac{n^2}{24} + O(n)$. \square

This is pretty nice. We can use calculus to get a decent lower bound of the problem. The calculation can even be done by hand. The hope to match the upper bound and lower bound is to try 11 intervals. This time we need a computer to help doing the calculation.

Example 3: $k = 11$,

We bound $T_{2,10}, T_{3,9}, T_{4,8}, T_{8,4}, T_{9,3}$ and $T_{10,2}$ by the area of each triangle which is $\frac{n^2}{242}$.

We bound $T_{i,12-i}$ by $S_{i,12-i}$, where $i = 1, 5, 6, 7, 11$.

We get eight optimal solutions to the maximum of Q . One of them is $[r_1, r_2, \dots, r_{11}] = [\frac{n}{11}, \frac{n}{11}, \frac{n}{11}, \frac{n}{11}, 0, 0, 0, 0, 0, 0, \frac{n}{11}]$.

This yields the maximum of Q as $\frac{9n^2}{22}$ which gives $M_\chi(n) \geq \frac{n^2}{22} + O(n)$. \square

YAY!! Since the lower bound matches the upper bound, the problem is solved.

3. GENERALIZED PROBLEM, $x + ay = z$, $a \geq 2$

3.1 A Greedy Algorithm for Upper bounds

We will show how to find an upper bound for the minimum number, over all 2-colorings of $[1, n]$, of monochromatic triples that are solutions of $x + ay = z$, for a fixed integer $a \geq 2$. We will obtain this upper bound by using the Greedy Algorithm. The general idea is the same as in the previous section. We again call the colors red and blue.

First

We paint the first interval of length k red. We will have $\frac{k^2}{2a}$ monochromatic triples as solutions of $x + ay = z$.

Second

We paint the second interval blue. We want to find the length of the interval (with this color) so that the overall number of monochromatic triples is minimum.

Let the length of this interval be $(a + j)k$ (here j is the number we want to find).

The total number of monochromatic triples on the whole interval is now $\frac{k^2}{2a} + \frac{j^2k^2}{2a} = \frac{(1+j^2)k^2}{2a}$.

The total length is $N = k + (a + j)k = (1 + a + j)k$.

So the total number of monochromatic triples in terms of N is

$$\frac{(1+j^2)(\frac{N}{1+a+j})^2}{2a} = \frac{(1+j^2)}{(1+a+j)^2} \frac{N^2}{2a}.$$

To find the minimum, we use calculus to get $j = \frac{1}{a+1}$. The total number of monochromatic Schur triples is then $\frac{N^2}{2a(a^2+2a+2)}$.

So far so good. We have a coloring that paints the first k integers red, followed by painting the next $(a + \frac{1}{a+1})k$ integers blue.

Third

Now we try to stick red at the end of the interval, and try to lower the overall number of triples. Say the length of this interval is jk , where j is the number we want to find.

The total number of monochromatic Schur triples on the whole interval is $\frac{k^2}{2a} + \frac{k^2}{2a(a+1)^2} + \frac{j^2k^2}{2a} = \frac{((a+1)^2+1+(a+1)^2j^2)k^2}{2a(a+1)^2}$.

The total length is $N = k + (a + \frac{1}{a+1})k + jk = (1 + a + \frac{1}{a+1} + j)k$.

So the total number of monochromatic Schur triples in term of N is $\frac{((a+1)^2+1+(a+1)^2j^2)}{2a(a+1)^2} (\frac{N}{(1+a+\frac{1}{a+1}+j)})^2 = \frac{((a+1)^2+1+(a+1)^2j^2)N^2}{2a((a+1)^2+1+(a+1)j)^2}$.

To find the minimum, we again use calculus to get $j = \frac{1}{a+1}$. The total number of monochromatic triples is $\frac{N^2}{2a(a^2+2a+3)}$. The coloring for the whole interval is a red interval of length equal to k , a blue interval of length equal to $(a + \frac{1}{a+1})k$ and another red interval of length equal to $\frac{1}{a+1}k$. k is such that the sum of these intervals is N , i.e. $k = \frac{N}{(1+a+\frac{2}{a+1})}$.

Fourth

We try to lower the bound even further by having a blue interval of length, say, jk at the end of the previous interval. But now we get that the minimizing j is negative. So we stop.

As a conclusion, the optimal coloring is proportional to $[1, a + \frac{1}{a+1}, \frac{1}{a+1}]$, with colors $[R, B, R]$ yielding that indeed the minimal number is $\frac{N^2}{2a(a^2+2a+3)}$.

3.2 Lower bounds

We will use a similar technique for the lower bound of the original problem. We find an upper bound for non-monochromatic triples in $[1, n]$. This gives a lower bound of the monochromatic triples.

We use the notation (R, B) and (B, R) for the non-monochromatic pair (x, y) .

Definition:

Let $|R|$ be the number of red points in $[1, n]$.

Let $|B|$ be the number of blue points in $[1, n]$.

Lemma 1) $|\{(R, B), (B, R) \mid y > x, y - x \text{ is divisible by } a\}| \leq \frac{|R||B|}{a}$.

Proof:

Let $|r_i|$ = number of red points at position m in $[1, n]$ where $m \equiv i \pmod{a}$.

Let $|b_i|$ = number of blue points at position m in $[1, n]$ where $m \equiv i \pmod{a}$.

We remark $r_i + b_i = \frac{n}{a}$, $1 \leq i \leq a$ and $\sum_{i=1}^a r_i = |R|$.

$$\begin{aligned} & |\{(R, B), (B, R) \mid y > x, y - x \text{ is divisible by } a\}| - \frac{|R||B|}{a} \\ &= \sum_{i=1}^a r_i b_i - \frac{|R||B|}{a} \\ &= \sum_{i=1}^a r_i \left(\frac{n}{a} - r_i \right) - \frac{(\sum_{i=1}^a r_i)(n - \sum_{i=1}^a r_i)}{a} \\ &= - \sum_{i=1}^a r_i^2 + \frac{(\sum_{i=1}^a r_i)^2}{a} \\ &\leq 0, \text{ by the Cauchy-Schwarz inequality.} \end{aligned}$$

Moreover, equality holds when $r_1 = r_2 = \dots = r_a$. \square

Let Q_a be two times the number of non-monochromatic triples of solutions of $x + ay = z$ in a 2-coloring of $[1, n]$.

Lemma 2) $Q_a \leq \frac{|R||B|}{a} + |\{(R, B), (B, R) \mid y - ax \geq 0\}| + |\{(R, B), (B, R) \mid y + ax \leq n\}|$.

Proof:


$$\begin{aligned} Q_a &= |\{\text{the non-monochromatic pair } (x, y) \mid \\ &\quad y > x \text{ and } y - x \text{ is divisible by } a\}| \\ &\quad + |\{\text{the non-monochromatic pair } (x, y) \mid y - ax \geq 0\}| \\ &\quad + |\{\text{the non-monochromatic pair } (x, y) \mid y + ax \leq n\}| \\ &\leq \frac{|R||B|}{a} + |\{(R, B), (B, R) \mid y - ax \geq 0\}| + |\{(R, B), (B, R) \mid y + ax \leq n\}| \\ &\quad, \text{ by lemma 1 } \square. \end{aligned}$$

When the points on the x -axis and the y -axis are painted with either color red or blue.

$|\{(R, B), (B, R) \mid y + ax \leq n\}|$ is the number of non-monochromatic coordinate pairs inside the triangle 1.

Similarly $|\{(R, B), (B, R) \mid y - ax \geq 0\}|$ is the number of non-monochromatic

coordinate pairs inside the triangle 2.



Divide the interval $[1, n]$ into k consecutive intervals.

Let r_i be the number of red points in the interval I_i .

Let b_i be the number of blue points in the interval I_i .

Let $S_{i,j}$ be the number of non-monochromatic pairs in the square of $I_i \times I_j$.

Let $T_{i,j}$ be the number of non-monochromatic pairs in the intersection of triangle and the square $I_i \times I_j$.

Note: $r_i + b_i = \frac{n}{k}$.

Theorem 3) $Q_2 \leq \frac{57n^2}{121} + O(n)$.

We find an upper bound of Q_2 by using calculus on the equation from the previous lemma.

The main part of calculating Q_2 is to compute the maximum number of non-monochromatic pairs in triangle 1 and triangle 2 in the pictures above. We partition an interval $[1, n]$ into k smaller intervals with equal length called them I_1, \dots, I_k . However there are $I_i \times I_j$ for some i, j that intersect the triangle only partly. We denote them $T_{i,j}$.

For each $T_{i,j}$ we have two ways to bound it

- 1) $T_{i,j} \leq \text{area of the intersection of triangle and the square } I_i \times I_j$.
- 2) $T_{i,j} \leq S_{i,j} = r_i b_j + r_j b_i$.

In this case, we use 11 intervals, $k = 11$.

In triangle 1, we bound $T_{1,10}, T_{2,9}, T_{2,8}, T_{3,7}, T_{3,6}, T_{5,3}, T_{5,2}$ and $T_{6,1}$ by the area of each intersecting triangle. We bound $T_{1,11}, T_{4,5}$ and $T_{4,4}$ by $S_{i,j}$.

In triangle 2, we bound $T_{2,4}, T_{3,5}, T_{3,6}$ and $T_{6,11}$ by the area of each intersecting triangle.

We bound $T_{1,1}, T_{1,2}, T_{2,3}, T_{4,7}, T_{4,8}, T_{5,9}$ and $T_{5,10}$ by $S_{i,j}$.

We then run the Maple program. We get four optimal solutions to the maximum of Q_2 . Two of them are

$$[r_1, r_2, \dots, r_{11}] = [\frac{n}{11}, \frac{n}{11}, 0, \frac{n}{11}, 0, 0, 0, 0, 0, \frac{n}{11}, 0] \text{ and } [\frac{n}{11}, \frac{n}{11}, 0, 0, 0, 0, 0, 0, 0, \frac{n}{11}, \frac{n}{11}, \frac{n}{11}].$$

The other two are the switching colors of the first two.

This yields the maximum of Q_2 as $\frac{57n^2}{121} + O(n)$. \square

Definition:

Let $M_{\chi,a}(n)$ be the number of monochromatic triples of solutions of $x + ay = z$ for a 2-coloring of $[1, n]$.

Corollary 4) $M_{\chi,2}(n) \geq \frac{7n^2}{484} + O(n)$.

Proof:

$$\begin{aligned} \text{The total number of triples} &= |\text{monochromatic triples}| + |\text{non-monochromatic triples}| \\ &= M_{\chi,a}(n) + \frac{Q_a}{2}. \end{aligned}$$

Since the total number of triples is $\frac{n^2}{2a} + O(n)$, we have $M_{\chi,a}(n) \geq \frac{n^2}{2a} - \frac{Q_a}{2} + O(n)$.

The lower bound, we found, of $M_{\chi,2}(n)$ follows from the upper bound of Q_2 from Theorem 3. \square

Note:

1) For $a = 3$, we found, $M_{\chi,3}(n) \geq \frac{n^2}{2268} + O(n)$. We ran the calculus program on 9 intervals with a particular upper bound of $T_{i,j}$.

2) For case $a \geq 4$, we could not find a positive lower bound for $M_{\chi,a}(n)$ yet. One of the reasons is that the upper bound of $M_{\chi,a}(n)$ is very small.

4. THE MINIMUM NUMBER, OVER ALL r -COLORING OF $[1, n]$, OF
MONOCHROMATIC SCHUR TRIPLES

4.1 A Greedy Algorithm for The Upper bound

The method to obtain the upper bounds in this section is similar to the one used in sections 2 and 3. In general we start with the first interval with color 1. Then we add interval 2 with color 2 in the optimal way. Then we add the third interval starting with color 1. If we get a positive solution, we move to the fourth interval. Otherwise we try with color 2. We keep going on in this fashion until there is no color that gives a positive solution.

Since there are many intervals involved in the computation, it is too much computation to do by hand. We wrote a computer program to help us compute the solutions for each r -coloring. The work was not so straightforward as we thought, since the details of counting triples and doing calculus at the same time are a bit tricky. But we made it at the end. We list the colorings up to $r = 5$, as examples, below. The program is available for download from the web site.

Definitions:

C = list of the coloring in order.

L = length of each interval (proportional to each other) corresponding to each color in C .

N = number of monochromatic Schur triple according to C and L .

$$\begin{array}{lll} r = 1, & C = [1], & L = [1], & N = \frac{n^2}{4} + O(n). \\ r = 2, & C = [1, 2, 1], & L = [1, \frac{3}{2}, \frac{1}{4}], & N = \frac{n^2}{22} + O(n). \\ r = 3, & C = [1, 2, 1, 3, 1, 2, 1], & L = [1, \frac{3}{2}, \frac{1}{4}, 3, \frac{1}{8}, \frac{487}{440}, \frac{47}{440}], & N = \frac{47n^2}{6238} + O(n) \\ & & & \sim \frac{n^2}{132.7234} + O(n). \end{array}$$

For $r \geq 4$, the lengths of the intervals are fractions with huge numerators and denominators. So we omit C and L here.

$$\begin{array}{l} r = 4, N = \frac{69631222699293042329481527n^2}{67076984091396704809405315398} + O(n) \sim \frac{n^2}{963.3176} + O(n). \\ r = 5, N \sim \frac{n^2}{7610.0730} + O(n). \end{array}$$

For $r = 6$, the lengths of the intervals are even larger fractions. This caused Maple to slow down. We waited for about 8 hours and we stopped. We did not get an answer.

However we were not really disappointed about this failure. The algorithm is more important.

4.2 Lower bounds

The method used to find a lower bound in the previous two sections could not be adapted for r -colorings, $r \geq 3$. We did not make any progress for a lower bound of r -coloring cases.

5. ABOUT THE PROGRAM

LowerBound(k, C)

input: the number of intervals k , list of types of upper bound C of $T_{i,k-i+1}$.

output: lower bound of $M_\chi(n)$, the upper bound of Q and the optimal solution of Q .

LowerBound2($k, C1, C2, a$)

input: the number of intervals k , list of types of upper bound $C1$ and $C2$ of $T_{i,k-i+1}$ and number a in equation $x + ay = z$.

output: lower bound of $M_{\chi,a}(n)$, the upper bound of Q_a and the optimal solution of Q_a .

minAllST(n, r)

input: length of intervals n , number of colors r .

output: the r -coloring of all the interval of length n that has the least number of monochromatic Schur triples.

Ord(C, L, n)

input: the list of coloring, the list of length corresponding to each color in C , symbol n .

output: the number of the monochromatic Schur triples of order n^2 .

Zeil(r)

input: number of color r .

output: the coloring with length of each coloring and also the total number of triples of order n^2 obtained from the Greedy Algorithm.

6. CONCLUSION

We have new upper bounds for generalized Schur triples $x + ay = z, a \geq 2$, in the 2-coloring case. We also have new upper bounds for Schur triples $x + y = z$, for r -colorings, $r \geq 3$ that considerably improve those of [2]. But we failed to match the lower and upper bounds

for these two problems. There is a possibility that other arguments in other papers [5], [1] and [4] for the lower bound used in the original problem can be adapted for the r -coloring problem. But the details of such an argument seem complicated. We believe these upper bounds are actually optimal. There might even be a beautiful simple way to solve it, but we failed to find it this time. We leave them as conjectures.

Conjectures:

- 1) The (asymptotic) number of minimum monochromatic triples of the form $\{x, y, x+ay\}$, $a \geq 2$ of 2-colorings of $[1, n]$, are $\frac{n^2}{2a(a^2+2a+3)} + O(n)$.
- 2) The (asymptotic) number of minimum Schur triples of r -colorings of $[1, n]$, $r \geq 3$, are the same as the upper bounds obtained from the Greedy Algorithm.

7. ACKNOWLEDGEMENT

I want to thank Bruce M. Landman and Aaron Robertson for their beautiful book that revealed Ramsey theory to all of us. I really enjoyed reading this book. I want to thank my advisor, Doron Zeilberger, for teaching me to do symbolic programming and making the graduate study years so much fun. I also want to thank Yoni Berkowitz for helpful discussions. Also thanks to my twin brother, Thotsaphon Thanatipanonda, for helping with the programming at the beginning phase of solving these problems.

REFERENCES

- [1] Boris Datskovsky, *On the number of monochromatic Schur triples*, Advances in Applied Math. 31(2003), 193-198.
- [2] Bruce M. Landman and Aaron Robertson, *Ramsey Theory on the Integers*, AMS, 2004.
- [3] Aaron Robertson, Pablo Parrilo and Dan Saracino, *On the Asymptotic Minimum Number of Monochromatic 3-Term Arithmetic Progressions*, JCTA 115 (2008), 185-192.
- [4] Aaron Robertson and Doron Zeilberger, *A 2-coloring of $[1, n]$ can have $\frac{n^2}{22} + O(n)$ monochromatic Schur triples, but not less!*, Electronic J. Combinatorics, 1998.

- [5] Tomasz Schoen, *The number of monochromatic Schur triples*, European J. Combinatorics 20(1999), 855-866.

DEPARTMENT OF MATHEMATICS

HILL CENTER-BUSCH CAMPUS

RUTGERS UNIVERSITY

110 FRELINGHUYSEN RD

PISCATAWAY, NJ 08854-8019

USA

E-mail address: `thot@math.rutgers.edu`