

Dr. Z's Math151 Handout #3.6 [Trigonometric Functions]

By Doron Zeilberger

Problem Type 3.6.1 : Differentiate an expression involving products and/or quotients of expressions containing *trig* functions.

Example Problem 3.6.1: Differentiate

$$y = \frac{1 + \cos x}{x + \sin x} .$$

Steps

1. Use the product or quotient rule to differentiate the given expression, remembering that $(\sin x)' = \cos x$, $(\cos x)' = -\sin x$, $(\tan x)' = \sec^2 x$. You don't really need to remember the more obscure formulas $(\csc x)' = -\csc x \cot x$, $(\sec x)' = \sec x \tan x$, $(\cot x)' = -\csc^2 x$, since you can always replace $\csc x$ by $1/\sin x$, etc.

2. Use algebra, and possibly trig identities, the most important one being

$$\sin^2 + \cos^2 = 1 ,$$

to simplify.

Example

1.

$$\begin{aligned} & \left(\frac{1 + \cos x}{x + \sin x} \right)' = \\ & \frac{(1 + \cos x)'(x + \sin x) - (1 + \cos x)(x + \sin x)'}{(x + \sin x)^2} = \\ & \frac{(-\sin x)(x + \sin x) - (1 + \cos x)(1 + \cos x)}{(x + \sin x)^2} . \end{aligned}$$

2.

$$\begin{aligned} & \frac{(-x \sin x - \sin^2 x) - (1 + 2 \cos x + \cos^2 x)}{(x + \sin x)^2} = \\ & \frac{-1 - x \sin x - 2 \cos x - (\sin^2 x + \cos^2 x)}{(x + \sin x)^2} = \\ & \frac{-1 - x \sin x - 2 \cos x - (1)}{(x + \sin x)^2} = \\ & \frac{-2 - x \sin x - 2 \cos x}{(x + \sin x)^2} . \end{aligned}$$

Ans.: $\frac{-2 - x \sin x - 2 \cos x}{(x + \sin x)^2}$.