


AUTHOR QUERY FORM

	Journal: YAAMA Article Number: 1412	Please e-mail or fax your responses and any corrections to: E-mail: corrections.essd@elsevier.vtex.lt Fax: +1 61 9699 6735
---	--	--

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the '[Q](#)' link to go to the location in the proof.

Location in article	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof
Q1	Please complete and update the reference(s) given here (preferably with a DOI if the publication data are not known): [3]. (p. 6/ line 52)
Q2	Please supply the year of publication. (Ref. [2]) (p. 6/ line 51)




ELSEVIER

Contents lists available at ScienceDirect

Advances in Applied Mathematics

www.elsevier.com/locate/yaama



The computational challenge of enumerating high-dimensional rook walks

Manuel Kauers^{a,*}, Doron Zeilberger^{b,2}

^a Research Institute for Symbolic Computation, J. Kepler University Linz, Austria

^b Mathematics Department, Rutgers University (New Brunswick), Piscataway, NJ, USA

ARTICLE INFO

Article history:

Received 10 March 2011

Accepted 14 March 2011

Available online xxxx

MSC:

05A15

33F10

68W30

Keywords:

Lattice walks

Computer algebra

Automated guessing

ABSTRACT

We present guesses, based on intensive computer algebra calculations, for recurrence equations of the sequences enumerating rook walks in up to twelve dimensions ending on the main diagonal. Computer proofs can in principle be constructed for all of them. For the moment, however, these computations are feasible only for low dimensions. We pose it as a challenge to develop algorithms which can also certify the correctness of the equations we found for the higher dimensions.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Consider a rook placed on the lower left corner $(0, 0)$ of a chess board. On how many paths can the rook reach the upper right corner (n, n) if in a single step it may move an arbitrary number of fields upwards or to the right (but not downwards or to the left)? For rectangular chess boards of size $n \times m$, the number of paths is given by the coefficient $a_{n,m}$ in the rational series expansion

$$\sum_{n,m=0}^{\infty} a_{n,m} x^n y^m = \frac{1}{1 - \frac{x}{1-x} - \frac{y}{1-y}}.$$

* Corresponding author.

E-mail addresses: mkauers@risc.jku.at (M. Kauers), zeilberg@math.rutgers.edu (D. Zeilberger).

¹ Partially supported by the Austrian FWF grant Y464-N18.

² Partially supported by the USA National Science Foundation.

The case of square chess boards is consequently the *diagonal series* of this rational function, which happens to be

$$\sum_{n=0}^{\infty} a_{n,n} x^n = \frac{1}{2} + \frac{1-x}{2\sqrt{1-10x+9x^2}}.$$

From here, all sorts of information about the numbers $a_{n,n}$ can be easily extracted by means of computer algebra, for instance the initial terms

$$1, 2, 14, 106, 838, 6802, \dots \quad (\text{A051708}),$$

or the recurrence equation

$$(n+2)a_{n+2,n+2} - (10n+14)a_{n+1,n+1} + 9na_{n,n} = 0 \quad (n \geq 0),$$

or the asymptotic formula $a_{n,n} \sim \sqrt{\frac{2}{\pi n}} 3^{2n-1}$ ($n \rightarrow \infty$). Computer algebra can also find the algebraic representation of the diagonal series given the bivariate rational series as input, so there is altogether no need to do any calculation by hand.

At a marvelous meeting at Nankai University in August 2010 on the occasion of the second author's 60th birthday, Frédéric Chyzak reported that he and his colleagues had succeeded in doing the analogous computation for 3D [3], i.e., they determined the number of paths a rook can take on a 3D chessboard from $(0, 0, 0)$ to (n, n, n) moving in each step an arbitrary positive integer distance into one of the three directions, i.e., moving either by $(i, 0, 0)$ or by $(0, i, 0)$ or by $(0, 0, i)$ for some positive integer i . Denoting now the number of this kind of walks by a_n , Martin Erickson, Suren Fernando, and Khang Tran [4] computed the initial terms

$$1, 6, 222, 9918, 486924, 25267236, \dots \quad (\text{A144045}),$$

conjectured the recurrence equation

$$\begin{aligned} &2(2+n)(3+n)^2(53+35n)a_{n+3} \\ &- (2+n)(43362+63493n+30114n^2+4655n^3)a_{n+2} \\ &+ (1+n)(54864+100586n+59889n^2+11305n^3)a_{n+1} \\ &- 192n^2(1+n)(88+35n)a_n = 0 \quad (n \geq 0), \end{aligned}$$

and proved asymptotic formula $a_n \sim \frac{9\sqrt{3}}{40\pi n} 64^n$ ($n \rightarrow \infty$) using a powerful analytical method of Robin Pemantle and Mark C. Wilson [7]. Chyzak and colleagues obtained a fully rigorous proof of the recurrence, including so-called certificates which allow for an independent formal verification of the obtained results. While it is clear *in theory* that computer algebra is able to obtain this information, it is remarkable that it is possible to actually carry out these calculations *in practice*, because the 3D case requires far more computational power than the 2D case.

If we don't insist on a fully rigorous formal verification, the diagonal recurrence can be obtained with much less effort: it suffices to compute some 25 terms of the sequence and use *automated guessing* to find a recurrence which matches them. This is also how Erickson and colleagues came up with their conjecture. For recent developments and references to classical versions of automated guessing, see [2,5]. For the present paper we applied this technique to empirically find recurrence equations for rook paths in dimensions greater than three, and we pose it as a challenge to provide rigorous certificates for them. While at least for the very high dimensions this seems totally hopeless

for now, we do expect that the coming years (or decades?) will see not only faster and bigger computers but also more advanced algorithms which can certify our claims within a reasonable amount of computing time. At least we intend to encourage progress in the development of more efficient algorithms. We see no other reason to ask for a certification. The question cannot be whether our claims are correct—the empirical evidence is way too strong to leave any reasonable doubt about that. Nor can the question be whether there actually exist certificates for our claims—it is clear by theory that recurrences of diagonal sequences of multivariate rational series can always be certified. Nor can the question be whether a proof may provide some insight or understanding—certificates are usually just messy polynomials. The interesting questions instead are: how big are the certificates, what is the computational cost for constructing them, and which techniques can be used to minimize the computational cost.

2. An alternative route for turning our semi-rigorous proofs to full-fledged rigorous proofs

We know *a priori* that there is a recurrence, this follows from general holonomic nonsense. But by the work of Moa Apagodu and Zeilberger [1] one can derive *a priori* upper bounds for the promised recurrences. The recurrences for d -dimensional rook walks turn out (empirically) to have order d (for $2 \leq d \leq 12$). It is very possible that it won't be too hard to prove this sharp upper bound in general, or even a weaker—but still realistic—one. This may enable one to give a “soft” proof that the empirically “guessed” recurrences are indeed *rigorously* proved.

If one would be able to find (realistic!) *a priori* bounds for the degrees of the coefficients as well, then by plain *linear algebra* the “guessed” recurrences would be rigorously proved.

3. A short interlude: Why is this problem so important?

The harsh and/or sceptical critic may say: Who cares? Not that many people (or machines) play 12-dimensional chess, and even the vast majority of the many people that do play traditional 2D, 8×8 chess, couldn't care less about the number of ways a rook can walk.

But *everyone* owes money, and usually to many creditors! The number of ways a rook can walk, in the d -dimensional cubic lattice, from the origin to $[n, \dots, n]$ is also the number of ways of repaying all your creditors if you currently owe n dollars to each of d different creditors, and a single payment consists of paying any positive amount of dollars (up to the whole debt) to any one of your creditors. Now this is a **very** practical problem.

4. Fast computation of sufficiently many sequence terms

As the dimension increases, so does the order of the diagonal recurrence and the degree of the polynomials appearing in it. The larger a recurrence is in terms of order and degree, the more sequence terms are needed to recover it from sequence data. For instance, in dimension $d = 12$, we needed 1600 diagonal terms in order to find the recurrence. To modern guessing software (we used code written by the first author [6]), this is still a moderate problem size. Much harder than guessing the recurrence is the computation of sufficiently many terms on the diagonals, which are needed as input for the guesser. The naive way is to start from the rational function

$$\frac{p(x_1, \dots, x_d)}{q(x_1, \dots, x_d)} = \frac{1}{1 - \frac{x_1}{1-x_1} - \frac{x_2}{1-x_2} - \dots - \frac{x_d}{1-x_d}} = \sum_{n_1, \dots, n_d=0}^{\infty} a_{n_1, \dots, n_d} x_1^{n_1} \dots x_d^{n_d}.$$

Its denominator $q(x_1, \dots, x_d)$ gives rise to a multivariate linear recurrence with constant coefficients, which can be used to compute the a_{n_1, \dots, n_d} recursively. For example, for $d = 2$, the rational function

$$\frac{1}{1 - \frac{x}{1-x} - \frac{y}{1-y}} = \frac{(x-1)(y-1)}{1 - 2x - 2y + 3xy}$$

implies the recurrence

$$3a_{n,m} - 2a_{n+1,m} - 2a_{n,m+1} + a_{n+1,m+1} = 0.$$

Together with suitable boundary conditions, this allows the computation of $a_{n,m}$ for arbitrary n, m , and hence for $a_{n,n}$ for arbitrary n .

But this is very costly. In dimension d , in order to compute the n th diagonal term, the recurrence forces us to compute all terms a_{n_1, \dots, n_d} with $0 \leq n_1, \dots, n_d \leq n$, altogether more than n^d terms. If $n = 1000$, a computer won't mind doing this for $d = 2$, but for $d = 3$ it is already getting painful, and for $d > 3$ either the memory requirements will exceed the computer's capacity or the runtime will exceed the user's patience. Or both. For $d \geq 10$ the naive method will not even suffice for computing the first $n = 10$ diagonal terms within a reasonable amount of time.

Of course, once we know a linear recurrence for the diagonal, we can compute the terms on the diagonal very efficiently. But this is of little use: if we knew linear recurrences for the diagonals already, we would not need to compute them. Fortunately, there are other recurrence equations, which are both efficient and easy to find. For arbitrary dimension d , we have



$$\begin{aligned} 1!n_d a_{n_1, \dots, n_{d-2}, n_{d-1}, n_d} &= (n_{d-1} - 1)a_{n_1, \dots, n_{d-2}, n_{d-1}-1, n_{d-1}} \\ &\quad + (n_{d-1} + 1)a_{n_1, \dots, n_{d-2}, n_{d-1}+1, n_{d-1}} \\ &\quad + (2 - n_d)a_{n_1, \dots, n_{d-2}, n_{d-1}, n_{d-2}} \\ &\quad + (2n_d - 2n_{d-1} - 2)a_{n_1, \dots, n_{d-2}, n_{d-1}, n_{d-1}}. \end{aligned}$$

The recurrence was discovered by multivariate automated guessing, and, once found, is easily proved in general. The key feature of this recurrence is that it leaves the indices n_1, \dots, n_{d-2} fixed, increases n_{d-1} and decreases n_d . This special form breaks the exponential complexity. It is not difficult to show that computing the first n diagonal terms via this recurrence requires only $O(n^2 d^3)$ operations. For $d \leq 7$, this method was efficient enough to produce enough terms to obtain the recurrence for the diagonal.

For $d \geq 8$, an additional improvement was needed. Here instead of directly computing the terms on the main diagonal, we first used the previous method for computing the terms of the bivariate auxiliary sequence

$$b_{n,m} := a_{n, \dots, n, m},$$

up to $n, m \leq 200$ or so. Then we used a multivariate guesser to discover some bivariate recurrences in n and m for $b_{n,m}$ and used these to compute the diagonal terms $b_{n,n} = a_{n, \dots, n, n}$ for n as far as needed.

5. Recurrence equations

Most of the recurrences we found are too big to be reproduced here. We make them available online at

<http://www.risc.jku.at/people/mkauers/walks/>.

Here we only give a table with some statistics on their order, the maximal degree of their polynomial coefficients, and the length of the longest integer appearing in them, measured in decimal digits (dd). The computation of this data took several weeks on a machine with eight processors and 32Gb of memory.

dim	ord	deg	maxint	OEIS tag	comment
2	2	1	2dd	A051708	easy
3	3	4	6dd	A144045	Chyzak et al.'s result
4	4	9	12dd	A181749	
5	5	18	31dd	A181750	
6	6	31	51dd	A181751	
7	7	50	94dd	A181752	
8	8	75	149dd	A181754	
9	9	108	236dd	A181725	
10	10	149	306dd	A181726	
11	11	200	462dd	A181727	
12	12	261	609dd	A181728	

6. Queens

We have also computed recurrences for the analogous problem of *Queen* walks, but so far we were only able to go up to dimension 5. The relevant output can be found in the above-mentioned webpage.

7. Higher order asymptotics

The leading-term asymptotics for diagonals of rook walks has been derived by Ericson et al. [4] using the powerful analytic method of Pemantle and Wilson [7]. It turns out to be

$$\sqrt{\alpha_d}(n\pi)^{(1-d)/2}(d+1)^{dn} \quad (n \rightarrow \infty),$$

where α_d is given by

$$\alpha_d = \frac{d^{d+2}}{(d+2)^{d-1}(d+1)^2 2^{d-1}}.$$

This result matches well with the numbers produced by the recurrences we discovered. For instance, for $d = 12$ and $n = 500000$ we find

$$\frac{\sqrt{\alpha_d}(n\pi)^{(1-d)/2}(d+1)^{dn}}{A181728(n)} = 1.0000020411 \dots$$

And this not all. Thanks to the Maple package *AsyRec* available from

<http://www.math.rutgers.edu/~zeilberg/tokhniot/AsyRec>

(see [8]) one can very easily get higher-order asymptotics from the recurrences, using the Birkhoff-Trjitznisky method.

The order-10 asymptotic formulas for the sequences for $2 \leq d \leq 9$ can be gotten from

<http://www.math.rutgers.edu/~zeilberg/tokhniot/oRookAsymptotics>

which is based on the input file

<http://www.math.rutgers.edu/~zeilberg/tokhniot/inRookAsymptotics>

which uses *AsyRec* and of course, the recurrences obtained by the first-named author's computer.

A cross check with $d = 12$ and $n = 500000$ now yields the very convincing quotient

[6] M. Kauers, *Guessing handbook*, Technical report 09-07, RISC-Linz, 2009.

[7] R. Pemantle, M.C. Wilson, Asymptotics of multivariate sequences. II. Multiple points of the singular variety, *Combin. Probab. Comput.* 13 (2004) 735–761.

[8] D. Zeilberger *AsyRec*, A maple package for computing the asymptotics of solutions of linear recurrence equations with polynomial coefficients, *Personal Journal of Shalosh B. Ekhad and Doron Zeilberger*, April 04, 2008, <http://www.math.rutgers.edu/~zeilberg/pj.html>.

UNCORRECTED PROOF