

Day 1 — Required

1–1. Let \mathbf{A} be an $n \times n$ (real) symmetric matrix.

- (a) Prove that all eigenvalues of \mathbf{A} are real.
- (b) Prove that two eigenvectors x and y of \mathbf{A} with distinct eigenvalues λ_x and λ_y are orthogonal.
- (c) If x is an eigenvector of A , find an $(n - 1)$ -dimensional \mathbf{A} -invariant subspace V of \mathbb{R}^n such that $x \notin V$. Recall that a subspace V of \mathbb{R}^n is \mathbf{A} -invariant iff for all $v \in V$, $\mathbf{A}v \in V$.

1–2. Find all possible values of

$$\oint_{\Gamma} \frac{e^{2\pi z}}{(z-1)(z-i)^2} dz$$

where Γ ranges over the class of simple, closed, smooth curves with $\Gamma \subset \mathbb{C} \setminus \{1, i\}$.

1–3. Let $f \in L^1(-\infty, \infty)$. Show that the functions $f(x)(\sin x)^n$, $n = 1, 2, \dots$, are measurable and integrable on $(-\infty, \infty)$ and

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} f(x)(\sin x)^n dx = 0.$$

Day 1 — Select 3

- 1–4. Let \mathbb{F}_2 be the field with 2 elements, and let $G = GL(3, \mathbb{F}_2)$ be the general linear group of 3×3 invertible matrices over \mathbb{F}_2 . Show that G has an element of order 7, and find all possible minimal polynomials for elements of order 7 in G .
- 1–5. Let $\{f_n(z)\}$ be a sequence of holomorphic functions defined in the open unit disk $D = \{z : |z| < 1\}$. Assume $\|f_n\|_{L^2(D)} \leq 1$. Show that $\{f_n\}$ has a subsequence which converges uniformly on every compact subset of D .
- 1–6. Let $\{E_k\}_{k=1}^\infty$ be a collection of Lebesgue measurable subsets of \mathbb{R}^n with $\sum_{k=1}^\infty |E_k| < \infty$. Show that the set consisting of all points which belong to infinitely many of the E_k has measure 0.
- 1–7. Let $R = \mathbb{Z} \oplus 2x\mathbb{Z}[x]$ be the subring of the ring of polynomials $\mathbb{Z}[x]$ given by

$$R = \{f(x) = a_0 + 2a_1x + 2a_2x^2 + \cdots + 2a_nx^n : a_i \in \mathbb{Z}\}.$$

That is, the constant term of a polynomial in R is any integer, but the coefficients of positive powers of x are even integers. Show that R is not Noetherian.

- 1–8. Let $z = x + iy$ and $f(z) = x(x^2 - 3y^2) + iy(3x^2 + y^2)$. State whether each of the following is true or false and give a reason.
- (a) $f'(0)$ exists.
 - (b) f satisfies the Cauchy-Riemann equations at $z = 0$.
 - (c) f is analytic (holomorphic) in a neighborhood of $z = 0$.
- 1–9. Let $R > 0$ and let f be an infinitely differentiable real-valued function on \mathbb{R}^n with support in a ball B_R of radius R centered at the origin. Recall that the support of a function $f(x)$ is the closure of the set of all x in the domain of f for which $f(x) \neq 0$.

- (a) Show that $|f(x)| \leq c \int_{B_R} \frac{|\nabla f(y)|}{|x - y|^{n-1}} dy$ for some constant c which depends on n but not on f, x or B_R . *Hint: start by applying the fundamental theorem of calculus along rays issuing from x .*
- (b) Using the above, show that if $n < p < \infty$, then there is a constant $c(n, p)$ such that
- $$|f(x)| \leq c(n, p) R^{1-n/p} \left(\int_{B_R} |\nabla f(y)|^p dy \right)^{1/p}.$$

Day 2 — Required

2-1. Consider the system of equations

$$\begin{aligned}t^2 + x^3 + y^3 + z^3 &= 0 \\t + x^2 + y^2 + z^2 &= 2 \\t + x + y + z &= 0\end{aligned}$$

and the point P with coordinates $(t, x, y, z) = (0, -1, 1, 0)$.

- (a) Show that the above equations determine a differentiable curve which can be expressed as

$$x = x(t), \quad y = y(t), \quad z = z(t)$$

in some neighborhood of the point P . Here “determine a differentiable curve” means that there is a differentiable function F with $F(t) = (x(t), y(t), z(t))$ and $F(0) = (-1, 1, 0)$.

- (b) Determine a nonzero tangent vector to the curve in (2-1.a) at the point P .

2-2. (a) Show that a compact metric space is complete.

- (b) Two metrics d_1 and d_2 on a space X are said to be “equivalent” if they generate identical topologies on X . If d_1 and d_2 are equivalent on X , does completeness of (X, d_1) imply completeness of (X, d_2) ? Prove it or give a counterexample.

2-3. For $n \geq 2$ let A_n denote the alternating group of even permutations of the set $\{1, 2, \dots, n\}$. Let K be the subgroup of A_n fixing a given element, say the element 1. If H is any subgroup of A_n with $[A_n : H] = n$, show that there is an isomorphism φ of A_n with itself such that $\varphi(H) = K$.

Day 2 — Select 3

2-4. Let $\{f_n\}$ and f be Lebesgue measurable functions on \mathbb{R} , and let m denote Lebesgue measure on \mathbb{R} . Determine whether or not each of the following statements is true or false, giving a proof if it is true, a counterexample if false.

- (a) If $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ for a.e. x , then $f_n \rightarrow f$ in measure as $n \rightarrow \infty$.
- (b) If $\lim_{n \rightarrow \infty} \int_{\mathbb{R}} |f_n - f| dm = 0$, then $f_n \rightarrow f$ in measure as $n \rightarrow \infty$.
- (c) Let f_n be integrable for each n , and let $g(x) = \liminf_{n \rightarrow \infty} f_n(x)$. Then

$$\int g(x) dm \leq \liminf_{n \rightarrow \infty} \int f_n dm.$$

2-5. Find a conformal mapping of the sector

$$\frac{\pi}{6} < \arg z < \frac{\pi}{3}$$

of the complex plane onto the open disk $|z| < 1$. Explain your answer.

2-6. Let $\vec{u}(x, y, z)$ denote the vector field

$$\vec{u}(x, y, z) = \left(\frac{x}{(x^2 + y^2 + z^2)^{3/2}}, \frac{y}{(x^2 + y^2 + z^2)^{3/2}}, \frac{z}{(x^2 + y^2 + z^2)^{3/2}} \right)$$

on $\mathbb{R}^3 \setminus \{(0, 0, 0)\}$, and let \mathcal{D} denote the annular region

$$1 < x^2 + y^2 + (z - 2)^2 < 9.$$

- (a) Compute the divergence of \vec{u} .
 - (b) Evaluate $\int \int_{\partial \mathcal{D}} \vec{u} \cdot \vec{n} ds$, where ds denotes the element of surface area of $\partial \mathcal{D}$ and \vec{n} denotes the unit outer normal to $\partial \mathcal{D}$.
- 2-7. Suppose that $f_n(x)$ is a continuous real-valued function on $[a, b]$ for $n = 1, 2, \dots$, and that for each $x \in [a, b]$, $\{f_n(x)\}$ is a bounded sequence. Prove that there is a subinterval of $[a, b]$ where $\{f_n(x)\}$ is uniformly bounded.

2-8. Let R be the subring of \mathbb{C} generated by the integers \mathbb{Z} and $\sqrt{-5} = i\sqrt{5}$, that is,

$$R = \mathbb{Z}[\sqrt{-5}].$$

- (a) Show that the element $x = 2 + \sqrt{-5}$ is not a product of two nonunits in R , that is, if $2 + \sqrt{-5} = rs$ for some r and s in R , then either r or s has an inverse in R .
- (b) Show that the ring R/xR is not an integral domain, where $x = 2 + \sqrt{-5}$.
- (c) Show that, for m and n nonzero integers, $m + 0\sqrt{-5}$ and $n + 0\sqrt{-5}$ have a greatest common divisor in R .

2–9. The exponential of a complex matrix \mathbf{A} is the power series

$$\exp(\mathbf{A}) = \sum_{i=0}^{\infty} \frac{(\mathbf{A})^i}{i!}.$$

You may use the properties that if \mathbf{A} and \mathbf{B} are commuting matrices, then $\exp(\mathbf{A} + \mathbf{B}) = \exp(\mathbf{A}) \cdot \exp(\mathbf{B})$, but this in general fails if $\mathbf{AB} \neq \mathbf{BA}$.

- (a) Show that any matrix \mathbf{A} is similar to a matrix \mathbf{B} which is a sum $\mathbf{B} = \mathbf{D} + \mathbf{N}$ of two commuting matrices \mathbf{D} and \mathbf{N} with \mathbf{D} diagonal and \mathbf{N} nilpotent, that is, \mathbf{N} has some power equal to 0.
- (b) It is a fact that the exponential $\exp(\mathbf{A}t)$ has columns which form a basis for the solution space of the system of differential equations $d\vec{\mathbf{X}}/dt = \mathbf{A}\vec{\mathbf{X}}$. Use this to find a basis for the solution space of the system

$$\begin{bmatrix} dx_1/dt \\ dx_2/dt \\ dx_3/dt \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$