

Week 2 The Dehn invariant, the Cauchy-Binet formula
 Jacobson I:7.1-7.2, Jacobson II: 1.1-1.3 , 3.1, 3.7-3.9

1. Let A and B be abelian groups.
 - a) Let m be a positive integer and let $a \in A$ be divisible by m in the sense that $a = ma'$ for some $a' \in A$. Show that $a \otimes b = a' \otimes mb$ in $A \otimes B$.
 - b) Let m and a be as in part a). Let $b \in B$ be an element of an abelian group B such that $mb = 0$ in B . Show that the element $a \otimes b$ is the identity element of $A \otimes B$.
 - c) Let A be an abelian group such that A is divisible in the sense that $A = mA$ for all positive integers m . Suppose that B is a torsion abelian group (that is, all elements of B have finite order). Show that $A \otimes B = 0$. Find an example of a nontrivial abelian group A such that $A \otimes A = 0$.
2. Show that in $\mathbf{R} \otimes \mathbf{R} / \pi \mathbf{Z}$ we have that for any nonzero integer m that $l \otimes \alpha / m = l / m \otimes \alpha$. Show that given an element $z \in \mathbf{R} \otimes \mathbf{R} / \pi \mathbf{Z}$ there is a finite set S of real numbers (perhaps empty) such that $z = \sum_{s \in S} l_s \otimes s$ such that the union of S and π is a linearly independent set over the rational field. Show that z is zero in the tensor product if and only if all $l_s = 0$. Show that $\mathbf{R} \otimes \mathbf{R} / \pi \mathbf{Z}$ is a torsion free abelian group.
3. Use the formula which computes the $k \times k$ minors of the product of a matrix A of size $m \times n$ and a matrix B of size $n \times p$ in terms of $k \times k$ minors of A and B in the special case $m = p = 2$ to prove Lagrange's identity: given elements $a_1, \dots, a_n, b_1, \dots, b_n$ of a commutative ring R then

$$\left(\sum_{l=1}^n a_l^2 \right) \left(\sum_{l=1}^n b_l^2 \right) - \left(\sum_{l=1}^n a_l b_l \right)^2$$

is a sum of squares in R . (Note this generalizes the Cauchy-Schwarz inequality for the case R the real numbers).